

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
4 August 2005 (04.08.2005)

PCT

(10) International Publication Number
WO 2005/069998 A2

(51) International Patent Classification: Not classified

(21) International Application Number:
PCT/US2005/002229

(22) International Filing Date: 19 January 2005 (19.01.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/537,622 20 January 2004 (20.01.2004) US
60/616,537 6 October 2004 (06.10.2004) US

(71) Applicant (for all designated States except US):
BRIGHAM YOUNG UNIVERSITY TECHNOLOGY TRANSFER OFFICE [US/US]; A285 ASB,
P.O.Box 21231, Provo, UT 84602-1231 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ANDRUS, Merritt, B.** [US/US]; 186 North 980 East, Lindon, UT 84042 (US).
LIU, Jing [CN/US]; 45 Wymount Terrace, Provo, UT 84604 (US).

(74) Agent: **WIGHT, Christopher, L.**; Holland & Hart LLP, Suite 3200, 555 - 17th Street, P.O: Box 8749, Denver, CO 80201 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.


(54) Title: NOVEL SIRTUIN ACTIVATING COMPOUNDS AND METHODS FOR MAKING THE SAME



(I)

(57) Abstract: The present invention includes methods for preparing resveratrol, resveratrol esters and substituted and unsubstituted stilbenes of the formula given below; where each Y is -O or halogen, each Z is -O or halogen, each n and each m is independently the value of 0, 1, 2, 3, 4 or 5, each A and each B is independently selected from P_n, R or absent, each V and each W is independently selected from

P_n, straight or branched alkyl of from (2) to (6) carbon atoms and cycloalkyl of from (3) to (8) carbon atoms, alkoxy, phenyl, benzyl or halogen, R is independently selected from the group comprising alkyl with at least one carbon atom, aryl and aralkyl, P_n is an alcohol protecting group and diastereoisomers of the foregoing. The compounds are made from a multi-step process including a N-heterocyclic carbene-type ligand coupling in the presence of a base with benzyol halide and styrene coupling partners. These compounds show increased stability for use in the food, cosmetic and pharmaceutical industries.



WO 2005/069998 A2

NOVEL SIRTUIN ACTIVATING COMPOUNDS AND METHODS FOR MAKING THE SAME

TECHNICAL FIELD

5 The present invention relates to compounds that activate sirtuin enzymes, and more particularly to analog compounds of resveratrol.

BACKGROUND

10 The sirtuins are a class of NAD^+ -dependent protein deacetylase enzymes that regulate a wide variety of cellular activities that promote cell survival and extend lifespan in response to environmental stress. Sirtuins exert their effect by removing acetyl groups from certain target proteins in the presence of oxidized nicotinamide adenine dinucleotide (NAD^+). For example, the yeast sirtuin enzyme Sir2 (silent information regulator 2), originally identified for its role in silencing transcription of
15 DNA, has also been shown to promote cell survival in response to caloric restriction. Similarly, in *C. elegans*, the sirtuin enzyme SIR-2.1 has been shown to extend lifespan. In mammalian cells, the sirtuin enzyme SIRT1 (a homolog of the yeast Sir2 and *C. elegans* SIR-2.1 enzymes) deacetylates the tumor suppressor p53 to promote cell survival. Sirtuins therefore appear to be activated as part of a
20 beneficial cellular response to stress, resulting in cell survival and extended lifespan. Activators of sirtuins may therefore be beneficial in effecting fundamental cellular processes that protect cells from stress and prevent or treat age-related diseases, and lengthen healthy life.

 Resveratrol (3,5,4'-trihydroxy-*trans*-stilbene) is a polyphenol compound,
25 known to be the most potent activator of sirtuins. As a stilbene phytoalexin, resveratrol continues to receive increasing attention for its role in mitigation of numerous and diverse human pathological processes including inflammation, atherosclerosis, and carcinogenesis. Resveratrol is known for its activity as an antioxidant, cyclooxygenase inhibitor, lipid modifier, platelet aggregation inhibitor
30 and vasodilator, inhibitor of tumor initiation, promotion and progression, neuroprotector, and antiviral compound.

 Resveratrol is an especially abundant component of red wines produced from grapes grown in cooler climates where plants are under stress from heavy disease

pressure. Indeed, the consumption of red wine containing resveratrol is believed to be responsible for the surprisingly normal lifespan of the French, despite their heavy consumption of fatty foods that cause heart disease, a phenomenon referred to as the “French Paradox.” Resveratrol, however, is not an optimal sirtuin activator. Its practical use is also limited due to difficult isolation and stereo-selectivity of extracts obtained from plant sources. More significantly, resveratrol is highly unstable due to its potential for oxidation, resulting in the formation of unstable radicals and quinones, and requiring that the final isolation of the product be carried out from impure mixtures containing multiple components.

Conventional methods of synthesizing resveratrol and resveratrol derivatives are expensive, inefficient, and do not provide for selective hydroxyl substitution. Many synthetic routes rely on Wittig and Horner-Emmons couplings that give mixtures of olefin isomers while still requiring 7-8 procedural steps. In addition, several conventional synthetic methods use methyl or benzyl ether protecting groups that require the use of boron tribromide or other inconvenient reagents for removal of those protecting groups.

In order to address the above problems, the present invention provides a novel and efficient route to synthetic resveratrol and novel derivative compounds of resveratrol having significantly improved activity and stability. Furthermore, the present invention provides a novel and efficient route to synthetic stilbene compounds.


DISCLOSURE THE INVENTION

The present invention provides novel stilbene compounds, in particular sirtuin activator compounds, and more particular ester analog compounds of resveratrol. These compounds have significantly improved biological activity and can be stored over a long period without alteration. The present invention also provides novel processes for synthesis of novel stilbene compounds, in particular novel sirtuin activator compounds, and more particular decarbonylative Heck couplings to synthesize stilbene derivatives. The present invention also provides an optional stilbene olefin isomerization procedure.

The instant invention is a compound of Formula I where each Y and each Z is independently selected from -O (ethers), -O-C=O, -C=O-O (esters), -O-C=O-O


(carbonates), -O-C=O-NH, -O-C=O-NR, -NH-C=O-O, -NR-C=O-O (carbamates),
 -NH-C=O, -NR-C=O, -C=O-NH, -C=O-NR (primary and secondary amides) -NH,
 -NR (primary and secondary amines), -N (heterocyclic rings), -S (thiol ethers), and
 halogen, each n and each m is independently the value of 0, 1, 2, 3, 4 or 5, where
 5 each A and each B is independently selected from H, R or absent, where each V and
 each W is independently selected from H, straight or branched alkyl of from 1 to 6
 carbon atoms and cycloalkyl of from 3 to 8 carbon atoms, alkoxy, phenyl, benzyl or
 halogen, and where R is independently selected from the group comprising alkyl
 with at least one carbon atom, aryl and aralkyl, and includes diastereoisomers of the
 10 foregoing. In some embodiments R is alkyl with at least two carbon atoms. In some
 embodiments the sum of m and n is greater than or equal to one but less than nine.
 In other embodiments the sum of m and n is three or more but less than nine.

Formula I




Preferred compounds are those of Formula I wherein each Y and each Z is
 15 independently selected from -O, -O-C=O and halogen, each n and each m is
 independently the value of 0, 1, 2, 3, 4 or 5, and the sum of n and m is 1 or more, V
 and W are each H, each A and each B is independently selected from H, R or absent,
 and where R is independently selected from the group comprising alkyl with at least
 one carbon atom, aryl and aralkyl, and includes diastereoisomers of the foregoing.
 20 In some embodiments R is alkyl with at least two carbon atoms. In some
 embodiments the sum of m and n is greater than or equal to one but less than nine.
 In other embodiments the sum of m and n is three or more but less than nine.

In a particular embodiment, compounds of Formula 1B




25 can be prepared where Y is independently selected from NH and O, n is equal to 0 1,
 2, 3, 4 or 5, Z is independently selected from NH and O, m is equal to 0 1, 2, 3, 4 or
 5, each A and each B is independently selected from R, P₁, P₂ and P₃ and each R is

independently selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing and P_1 , P_2 and P_3 are protecting groups, comprising the step of coupling a benzoyl halide compound of Formula 1B2




5 where X is halogen, with styrene compound of Formula 1B3




in solution or in suspension of a first solvent, transition metal catalyst, N-heterocyclic carbene-type ligand and a first base. In other embodiments, each R are independently selected from alkyl with at least two carbon atoms.

10 In another aspect of the invention, the resveratrol (3,5,4'-trihydroxy-*trans*-stilbene) esters of Formula II are characterized in that they comprise at least one ester group of formula $-O-C(=O)-R$ at one or more of the 3, 5 and 4' positions, wherein R represents an alkyl with at least one carbon atom, aryl and aralkyl, and includes diastereoisomers of the foregoing. In some embodiments R is alkyl with at
15 least two carbon atoms. In some embodiments the sum of m and n is greater than or equal to one but less than nine. In other embodiments the sum of m and n is three or more but less than nine.

Formula II




In a particular embodiment, compounds of Formula 1A3



20


can be prepared where A_1 is selected from P_1 and $(CO)R_1$; A_2 is selected from P_2 and $(CO)R_2$; A_3 is selected from P_3 and $(CO)R_3$; where P_1 when present is a first alcohol protecting group, P_2 when present is a second alcohol protecting group, P_3

when present is a third alcohol protecting group, R₁, R₂ and R₃ when present are each independently selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing, comprising the step of coupling a benzoyl halide compound of Formula 1A4



5

where X is halogen, with a 4-substituted styrene compound of Formula 1A5




in solution or in suspension of a first solvent, transition metal catalyst, N-heterocyclic carbene-type ligand and a first base. In other embodiments, R₁, R₂ and R₃ when present are each independently selected from alkyl with at least two carbon atoms.

In some embodiments, R represents a saturated or unsaturated fatty acid group or radical. In the case of an unsaturated fatty acid, the double bonds are advantageously *cis*, which corresponds to the most frequent case found in the natural products. With products obtained more particularly by synthesis or hemisynthesis, the bonds are *trans*. Fatty acid derivatives which are suitable for the implementation of the invention include but are not limited to: butyric C4:O; valeric, C5:O; hexanoic, C6:O; sorbic, C6:2(n-2); lauric C12:O; palmitic C16:O; stearic, C18:O; oleic, C18:1(n-9); linoleic, C18:2(n-6); linolenic, C18:3(n-6); .alpha. linolenic, C18:3(n-3); arachidonic, C20:4(n-3); eicosapentaenoic C20:5(n-3); and docosahexanoic, C22:6(n-3). The C16 and more fatty acids are particularly appropriate as regards cosmetic uses. These fatty acids are extracted, for example, from microalgae.

In another embodiment, R represents an aryl group, the aryl group being more particularly phenyl, benzyl or styryl.

In yet another embodiment, R represents an aralkyl or aralkylene group, the alkyl or alkylene group being more particularly C₁ to C₈, in particular C₁ to C₄. In particular the aryl portion is phenyl, benzyl or styryl group.

In still another aspect of the invention, an improved process for the preparation of compounds of Formulas I and II involves a decarbonylative coupling of compounds of Formula 1C and Formula 1E in solution or in suspension of a solvent, transition metal catalyst, N-heterocyclic carbene-type ligand and base, where X is halogen, and each A and each B is independently selected from a protecting group and alkyl with at least one carbon atom, aryl and aralkyl, or absent when A or B is halogen.



In yet another aspect of the invention, an improved process for the preparation of resveratrol arises from the 1) acetoxylation of 3,5-dihydroxy benzoic acid with an acetoxy esterification agent in solution with a base; 2) halogenation in solution; 3) reaction with an acetoxy styrene in solution, transition metal catalyst, N-heterocyclic carbene-type ligand, and base; and, 4) hydrolysis in solution with an alkali metal hydroxide.

In another aspect of the invention, novel benzoic acid derivatives of Formula 1C are characterized in that each Y is independently selected from -O, -O-C=O, A is H or R, n is equal to the value of 0, 1, 2, 3, 4 or 5, X is any halogen, and R is independently selected from the group comprising alkyl with at least one carbon atom, aryl, and aralkyl. In some embodiments R is alkyl with at least two carbon atoms. In some embodiments the sum of m and n is greater than or equal to one but less than nine. In other embodiments the sum of m and n is three or more but less than nine.

In still another aspect of the invention, a novel process for preparing derivatives of benzoic acid halides of the Formula 1C are characterized in that said compounds are halogenated with a halogenating agent in a solvent and benzotriazole. Preferred halogenating agents are thionyl chloride and oxalyl chloride.

In yet another aspect of the invention, preferred compounds of Formula 1C are those where X is Cl.

In another aspect of the processes of the invention, the base used in the step for coupling an acid chloride with a styrene derivative is a non-coordinating amine base. Preferred non-coordinating amine bases are selected from the group comprising: N,N-dimethylbenzylamine, N-methylmorpholine,
5 dimethylaminopyridine. A more preferred non-coordinating amine base is N,N-dimethylbenzylamine.

In another aspect of the invention, an N-heterocyclic carbene-type ligand is employed in a decarbonylative Heck reaction. Preferred N-heterocyclic carbene-type ligands are selected from imidazolium, 1,3-disubstituted imidazolium (N,N'-
10 bis-substituted imidazolium), 1,3-disubstituted-4,5-dihydroimidazolium carbene-type ligands. More preferred N-heterocyclic carbene-type ligands are those selected from N,N'-bis-carbocycle imidazolium ligands. Even more preferred N-heterocyclic carbene-type ligands are those selected from N,N'-bis-(2,6-diisopropylphenyl) imidazolium chloride, N,N'-bis-(dimesityl) imidazolium chloride, N,N'-bis-
15 adamantylimidazolium chloride.

In still another aspect of the invention, a halogenating agent is used. Preferred halogenating agents are selected from the group comprising: thionyl chloride, thionyl bromide, thionyl iodide, oxalyl chloride, oxalyl bromide, oxalyl iodide, PCl₃, PBr₃, PI₃, POCl₃, POBr₃, and POI₃. More preferred halogenating
20 agents are thionyl chloride and oxalyl chloride.

In another aspect of the invention, an acylating agent is used. Preferred acylating agents are selected from the group comprising: ester anhydrides, mixed ester anhydrides, acid halides. A more preferred acylating agent is acetic anhydride.

In yet another aspect of the invention a process includes the use of a
25 protecting agent. Preferred protecting agents are those selected from the group comprising MOMCl, lev₂O, and chloroacetoxy chloride.

In another aspect of the invention, a hydrolysis step is performed using a reagent selected from alkaline earth hydroxides and alkoxides. Preferred hydrolysis reagents are lithium hydroxide, sodium hydroxide, potassium hydroxide, lithium
30 methoxide, sodium methoxide, lithium methoxide, and lithium ethoxide.

In another aspect of the invention includes the use of a transition metal catalyst. Preferred transition metal catalysts are palladium, rhodium, and ruthenium catalysts. More preferred transition metal catalysts are palladium catalysts.

Preferred palladium catalysts are selected from the group comprising: Pd(OAc)₂, PdCl₂, PdBr₂, palladium bis triphenylphosphine, and palladium tris triphenyl phosphine. A more preferred Pd II catalyst is Pd(OAc)₂.

In still another aspect of the invention, a deprotection reagent is used.

- 5 Preferred deprotection agents are those selected from the group comprising: sodium sulfite and sodium thiosulfite, aqueous pyridine at pH 6.7, NaI and TMSCl, and alkaline earth metal hydroxide and a polar aprotic solvent.

- 10 In yet another aspect of the invention, pharmaceutical compositions of the compounds of Formulas I and II comprising a therapeutically effective amount of a compound of Formulas I and II in admixture with a pharmaceutically acceptable carrier are described.

In another aspect of the invention, cosmetic compositions of the compounds of Formula I and II comprising a cosmetically effective amount of a compound of Formula I and II in admixture with a cosmetically acceptable carrier are described.

- 15 In still another aspect of the invention, nutritional compositions of the compounds of Formula I and II comprising a nutritionally effective amount of a compound of Formula I and II in admixture with an nutritionally acceptable carrier are described.

- 20 The compounds Formula I and II are sirtuin activator compounds that interact with mammalian enzymes including SIRT1 promoting cell survival.

In still another aspect of the invention, levulinic halides and anhydrides are used to protect phenolic hydroxyls.

BEST MODE(S) FOR CARRYING OUT THE INVENTION

- 25 In the compounds of Formulas I, II and 1-7 and as further used herein, the term "heteroatom" means nitrogen, oxygen, or sulfur. The term "halogen" as used herein means bromine, chlorine, fluorine, or iodine.

- 30 The term "cis" as used herein means the stereoisomerism about a carbon-carbon double bond, a monocyclic ring, a fused bicyclic ring, or a bridged bicyclic ring wherein the highest ranking substituent on each of the two carbons of relevance are on the same side, which substituent ranking is based on the sequence rules of the Cahn-Ingold-Prelog system (March, J., Advanced Organic Chemistry, 4th ed., 1992, New York: John Wiley & Sons, 1992:109, 127-133 and references cited therein).

The term “trans” means the stereoisomerism about a carbon-carbon double bond, a monocyclic ring, a fused bicyclic ring, or a bridged bicyclic ring wherein the highest ranking substituent on each of the two carbons of relevance are on opposite sides, which substituent ranking is based on the sequence rules of the Cahn-Ingold-Prelog system (March J., *supra.*, 1992; 109:127-133 and references cited therein).

The term “alkyl” as used herein means a straight or branched hydrocarbon radical or group having at least one carbon atom including but not limited to saturated C₁-C₆ such as: methyl, ethyl, 1-propyl and 2-propyl, 1-butyl, 2-butyl, 2-methyl-1-propyl, 1,1-dimethylethyl, 1-pentyl, 2-pentyl, 3-pentyl, 2-methyl-1-butyl, 3-methyl-1-butyl, 2,2-dimethylpropyl, 1-hexyl, 2-hexyl, 3-hexyl, 2-methyl-1-pentyl, 3-methyl-1-pentyl, 4-methyl-1-pentyl, 3,3-dimethyl-1-butyl, 3,3-dimethyl-2-butyl, 2-ethyl-1-butyl and the like; C₇-C₁₂ such as: 1-heptyl, 2-heptyl, 3-heptyl, 4-heptyl, 2-methyl-1-hexyl, 4-methyl-1-hexyl, 5-methyl-1-hexyl, 1-octyl, 2-octyl, 3-octyl, 4-octyl, 6-methyl-1-heptyl, 5,5-dimethyl-1-hexyl, 2-ethyl-1-hexyl, 2-methyl-1-heptyl, 2-propyl-1-pentyl, 1-nonyl, 2-nonyl, 2-ethyl-2-methyl-1-hexyl, 4-methyl-1-octyl, 3,5,5-trimethyl-1-hexyl, 1-decyl, 2-decyl, 4-ethyl-1-octanyl, 2-methyl-1-nonyl, 4-methyl-1-nonyl, 8-methyl-1-nonyl, 1-undecyl (1-hendecyl), 2-undecyl, 7-methyl-1-decyl, 1-dodecyl, 5-dodecyl, 2-butyl-1-octyl, 10-methyl-1-undecyl and the like; C₁₃-C₁₈ such as: 1-tridecyl, 4-methyl-1-dodecyl, 11-methyl-1-dodecyl, 1-butyldecyl, 11-methyl-1-tridecyl, 1-pentadecyl, 1-hexadecyl, 2-hexyl-1-decyl, 1-heptadecyl, 14-methyl-1-hexadecyl, 15-methyl-1-hexadecyl, 1-octadecyl, 16-methyl-1-heptadecyl and the like; C₁₉-C₃₂ such as 1-nonadecyl, 2-methyl-1-octadecyl, 10-methyl-1-octadecyl, 17-methyl-1-octadecyl, 2,6,10,14-tetramethylpentadecyl, 1-eicosyl (1-arachidinyl, 1-leicosanyl), 18-methyl-1-nonadecyl, 1-heneicosyl, 19-methyl-1-eicosyl and 1-docosyl (1-behenyl), 1-tricosyl, 1-tetracosyl, 1-pentacosyl, 1-hexacosyl, 1-heptacosyl, 1-octacosyl, 1-nonacosyl, 1-triacontyl, 2,6,10,15,19,23-hexamethyl-1-tetracosyl, 1-hentriaconsyl, 1-dotriacontyl and the like. Alkyl groups may be unsubstituted or substituted.

Alkyl groups having two or more carbons may optionally contain 1 or more sites of unsaturation, the groups being known as alkenyl groups or radicals and alkynyl groups or radicals. Alkenyl groups are analogous to alkyl groups which are saturated, but have at least one double bond (two adjacent sp² carbon atoms). Depending on the placement of a double bond and substituents, if any, the geometry

of the double bond may be *trans* (E), or is (Z). Similarly, alkynyl groups have at least one triple bond (two adjacent sp carbon atoms). Unsaturated alkenyl or alkynyl groups may have one or more double or triple bonds, respectively, or a mixture thereof. Like alkyl groups, unsaturated groups may be straight chain or branched.

- 5 Examples of alkenyls include vinyl, allyl, 2-methyl-2-propenyl, cis-2-butenyl, trans-2-butenyl, and acetyl, propene, 1-butene, 2-butene, 2-methylpropene, 1-pentene, 2-pentene, 2-methyl-1-butene, 2-methyl-2-butene, 3-methyl-1-butene, 1-hexene, 2-hexene, 3-hexene, 2,3-dimethyl-1-butene, 2,3-dimethyl-2-butene, 3,3-dimethyl-1-butene, 2,4-dimethyl-2-butene, 2-ethyl-1-butene, 2-methyl-1-pentene, 2-methyl-2-pentene, 3-methyl-1-pentene, 3-methyl-2-pentene, 4-methyl-1-pentene, 4-methyl-2-pentene, 1-heptene, 2-heptene, 3-heptene, 3,4-dimethyl-2-pentene, 4,4-dimethyl-2-pentene, 3-methyl-2-hexene, 3-methyl-3-hexene, 4-methyl-2-hexene, 2,3-dimethyl-1-pentene, 2,3-dimethyl-2-pentene, 2,4-dimethyl-1-pentene, 2,4-dimethyl-2-pentene, 3,3-dimethyl-1-pentene, 3,4-dimethyl-1-pentene, 4,4-dimethyl-1-pentene, 4,4-dimethyl-2-pentene, 3-ethyl-1-pentene, 3-ethyl-2-pentene, 2-methyl-1-hexene, 2-methyl-2-hexene, 3-methyl-1-hexene, 4-methyl-1-hexene, 5-methyl-1-hexene, 2,3,3-trimethyl-1-butene, 1-octene, 2-octene, 3-octene, 4-octene, 2,2-dimethyl-3-hexene, 2,3-dimethyl-2-hexene, 2,3-dimethyl-3-hexene, 3-ethyl-2-methyl-1-pentene, 3-ethyl-2-methyl-pent-2-ene, 2-isopropyl-1-pentene, 2-methyl-1-heptene, 2-methyl-2-heptene, 4-methyl-2-heptene, 2,3,4-trimethyl-2-pentene, 2,4,4-trimethyl-1-pentene, 2,4,4-trimethyl-2-pentene, 3,4,4-trimethyl-2-pentene, 1-nonene, 2-nonene, 3-nonene, 4-nonene, 2,2-dimethyl-3-heptene, 3,5,5-trimethyl-1-hexene, 1-decene, 4-decene, 5-decene, 3,7-dimethyl-1-octene, 2-methyl-1-nonene, 1-undecene, trisisobutylene, 2,2,4,6,6-pentamethyl-3-heptene, 1-dodecene, 2-methyl-1-undecene, 1-tridecene, 1,1-dimethylethylene, 1-tetradecene, 7-tetradecene, 1-pentadecene, 1-hexadecene, 1-heptadecene, 8-heptadecene, 1-octadecene, 1-nonadecene, 1-eicosene, 9-heneicosene, 1-docosene and the like.

- Examples of dialkenes include but are not limited to propadiene (allene), 1,3-butadiene, 1,3-pentadiene, 1,4-pentadiene, 2-methyl-1,3-butadiene (isoprene), 3-methyl-1,2-butadiene, 1,3-hexadiene, 1,4-hexadiene, 1,5-hexadiene, 2,4-hexadiene, 2,3-dimethyl-1,3-butadiene, 2-methyl-1,3-pentadiene, 2-methyl-1,4-pentadiene, 3-methyl-1,4-pentadiene, 4-methyl-1,3-pentadiene, 3-methyl-1,3-pentadiene 1-heptyne, 2-heptyne, 3-heptyne, 2,4-dimethyl-1,3-pentadiene, 2,4-dimethyl-2,3-

pentadiene, 1,6-heptadiene, 1,7-octadiene, 1,4-octadiene, 3-methyl-1,5-heptadiene, 2,5-dimethyl-1,5-hexadiene, 2,5-dimethyl-1,4-hexadiene, 1,8-nonadiene, 7-methyl-1,6-octadiene, 1,9-decadiene, 7-dimethyl-1,6-octadiene, 5,7-dimethyl-1,6-octadiene, 1,7-hexadecadiene and the like.

5 Examples of trialkenes include but are not limited to 5-methyl-1,3,6-heptatriene, 2,6-dimethyl-2,4,6-octatriene (neo-alloocimene), 1,5,9-decatriene, 2,6-dimethyl-2,4,6-octatriene, 3,7-dimethyl-1,3,6-octatriene, 7-methyl-3-methylene-1,6-octadiene, 3,7-dimethyl-1,3,6-octatriene, 1,4,9-decatriene, 1,3,5-undecatriene and the like. Examples of alkynyls include, but are not limited to 1-butyne, 2-butyne, 1-
10 pentyne, 2-pentyne, 4-methyl-pent-1-yne, 1-hexyne, 2-hexyne, 3-hexyne, 3,3-dimethyl-1-butyne, 1-heptyne, 2-heptyne, 3-heptyne, 5-methyl-1-hexyne, 1-octyne, 2-octyne, 3-octyne, 4-octyne, 1-nonyne, 1-decyne, 5-decyne and 1-dodecyne, 1-pentadecyne and the like. Alkenyl and Alkynyl groups may be unsubstituted or substituted.

15 As used herein, alkyl also includes mixed alkenyl and alkynyl groups. An unsaturated hydrocarbon may thus include subunits of double bonds and subunits of triple bonds. Examples of these mixed alkenyl and alkynyl groups include 2-methyl-1-buten-3-yne, 2-methyl-1-hexen-3-yne and the like. Mixed alkenyl and alkynyl groups may be unsubstituted or substituted.

20 Alkyl also includes groups having three or more carbons that contain 1 or more sites of unsaturation, that group being known as cycloalkyl groups or radicals.

 The term "cycloalkyl" as used herein means a monocyclic or polycyclic hydrocarbyl group. Illustrative examples of a cycloalkyl group or radical include cyclopropyl, cycloheptyl, cyclooctyl, cyclodecyl, cyclobutyl, adamantyl, norpinanyl,
25 decalinyl, norbornyl, cyclohexyl, and cyclopentyl. Cycloalkyl groups may be unsubstituted or substituted. Also included are rings in which 1 to 3 heteroatoms replace carbons. Such groups are termed "heterocyclyl", which means a cycloalkyl group also bearing at least one heteroatom selected from O, S, or N, examples being oxiranyl, pyrrolidinyl, piperidyl, tetrahydropyran, and morpholine.

30 The term "aryl" as used herein means an aromatic carbocyclic ring having from 6 to 14 carbon atoms. Illustrative examples of an aryl group or radical include phenyl, 1-naphthyl, 2-naphthyl, 1-anthryl, 2-anthryl, 9-anthryl, 1-phenanthryl, 2-phenanthryl, 3-phenanthryl, 4-phenanthryl, 5-phenanthryl, and the like; including

fused ring systems with rings that have less than 6 carbons such as 1-acenaphthyl, 3-acenaphthyl, 4-acenaphthyl, 5-acenaphthyl, 1-azulyl, 2-azulyl, 4-azulyl, 5-azulyl, 6-azulyl and the like. Aryl groups may be unsubstituted or substituted.

The term "aryl" also includes heteroaryls. The term "heteroaryl" means an
5 unsaturated monocyclic group or radical of 5 or 6 atoms, an unsaturated fused bicyclic group or radical of from 8 to 10 atoms, or an unsaturated fused tricyclic group or radical of from 11 to 14 atoms, the cyclic groups having 1 or 2 heteroatoms independently selected from O, N, or S. Illustrative examples of monocyclic heteroaryl include 2-or 3-thienyl, 2-or 3-furanyl, 1-, 2-, or 3-pyrrolyl, 1-, 2-, or 4-
10 imidazolyl, 1-, 3-, or 4-pyrazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 3-, 4-, or 5-isoxazolyl, 3-, 4-, or 5-isothiazolyl, 2-, 3-, or 4-pyridinyl, 3-or 4-pyridazinyl, 2-or 3-pyrazinyl, and 2-, 4-, or 5-pyrimidinyl. Illustrative examples of bicyclic heteroaryl include 2-, 3-, 4-, 5-, 6-, 7-, or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, or 8-isoquinolinyl, 1-, 2-, 3-, 4-, 5-, 6-, or 7-indolyl, 2-, 3-, 4-, 5-, 6-, or 7-
15 benzo[b]thienyl, 2-, 4-, 5-, 6-, or 7-benzofuran, 2-, 4-, 5-, 6-, or 7-benzoxazolyl, 2-, 4-, 5-, 6-, or 7-benzothiazolyl, and 1-, 2-, 3-, 4-, 5-, 6-, or 7-benzimidazolyl. Illustrative examples of tricyclic heteroaryl include 1-, 2-, 3-, or 4-dibenzofuranyl, 1-, 2-, 3-, or 4-dibenzothienyl, and 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8-, or 9-(1,2,3,4-tetrahydroacridinyl). Heteroaryl groups may be unsubstituted or substituted.

20 As used above, a fused bicyclic group or radical is a group wherein two ring systems share two and only two atoms. As used above, a fused tricyclic group or radical is a group wherein three ring systems share four and only four atoms.

The term "aralkyl" as used herein means an aryl-alkyl group or radical wherein aryl and alkyl have the meanings as defined above including where alkyl
25 includes alkenyl, alkynyl and cycloalkyl and where aryl includes heteroaryl. Illustrative examples of an arylalkyl group or radical include benzyl, 4-fluorophenylmethyl, 2-phenylethyl, 3-phenylpropyl, 4-phenylbutyl, 3-methyl-3-phenylpropyl, 1-naphthylmethyl, 1-naphthylethyl, 3-(1-naphthyl)-propyl, 4-(1-naphthyl)-butyl, 4-(2-naphthyl)-butyl, 4-phenylheptyl, and 12-(2-hydroxyphenyl)-
30 dodec-3-yl.

The term "alcohol protecting group" as used herein means a conventional alcohol protecting group known to those skilled in the art such as is used to protect a hydroxy group. Such alcohol protecting groups are described in T. W. Greene,

Protective Groups in Organic Synthesis, John Wiley & Sons, New York, N.Y., 1991, 2nd Edition, which is hereby incorporated by reference and include for example, esters such as formyl, (C₁-C₁₀) alkanoyl optionally mono-, di- or tri-substituted with (C₁-C₆) alkoxy, halo, aryl, aryloxy or haloaryloxy; aroyl optionally
5 mono-, di- or tri-substituted on carbon with halo, (C₁-C₆) alkyl, (C₁-C₆) alkoxy wherein aryl is phenyl, 2-furyl etc; carbonates; sulfonates; and ethers such as benzyl, paramethoxybenzyl, methoxymethyl, etc. The lev protecting group is discussed by van Boom, J.H.; Burgers, P.M.J. Tetrahedron Letters, 1976, 4875.

The term "N-heterocyclic carbene-type ligand" as used herein means N-
10 heterocyclic carbene-type ligands generated from their corresponding salts. Preferred N-heterocyclic carbene-type ligands are those which include bulky groups that shield the carbene from dimerizing. Preferred N-heterocyclic carbene-type ligands include imidazolium, 1,3-disubstituted imidazolium (N,N'-bis-substituted imidazolium), 1,3-disubstituted-4,5-dihydroimidazolium carbene-type ligands. More
15 preferred 1,3-disubstituted imidazolium carbene-type ligands include N,N'-bis-carbocycle imidazolium ligands. In the N,N'-bis-carbocycle imidazolium ligands, the term "carbocycle" (and variations thereof such as "carbocyclic" or "carbocyclyl") as used herein broadly refers to substituted and unsubstituted, saturated or unsaturated, mono, bi, and tricyclic ring systems in which the rings are
20 independent or fused and in which each ring is saturated or unsaturated. Examples of an unsaturated substituted monocyclic N,N'-bis-carbocycle-imidazolium salts are: N,N'-bis-(2,6-diisopropylphenyl) imidazolium chloride (also called 1,3-di-(2,6-diisopropylphenyl) imidazolium chloride) and N,N'-bis-(dimesityl) imidazolium chloride (also called 1,3-di-(2,4,6-trimethylphenyl)imidazolium chloride). An
25 example of a saturated unsubstituted tricyclic N,N'-di-carbocycleimidazolium salt is N,N'-bis-adamantylimidazolium chloride.

The term "transition metal catalyst" as used herein means any transition metal useful in forming carbon-carbon bonds including salts, carbonyl compounds, chelates, or complexes with ligands having trivalent donor groups of the metals in
30 Group VIII or copper. A Group VIII metal includes Fe, Co, Ni, Ru, Rh, Pd, Os, Ir, or Pt. Complex catalysts are those with ligands having trivalent donor atoms and are comprised of a Group VIII metal complexed by one or more ligands. These complexes are formed by the reaction of a Group VIII metal compound and a ligand

having a trivalent donor atom. Such trivalent donor atoms include phosphorus, nitrogen, arsenic, antimony and bismuth. These types of complexes are well known to those of ordinary skill in the art and most commonly involve phosphorus-type ligands. (see R. F. Heck, *Palladium Reagents in Organic Syntheses*, Academic Press, N.Y., N.Y., 1985, pages 1-7; S. G. Davies, *Organotransition Metal Chemistry Applications to Organic Syntheses*, Pergamon Press, N.Y., N.Y., 1985, pages 13-17). Examples of the most common phosphorus-type ligands include phosphines of which the preferred are triphenylphosphine, trimethylphosphine, methyldiphenylphosphine, dimethylphenylphosphine, dicyclohexylphenylphosphine, diphenylenephénylphosphine, tri-p-tolylphosphine, tri-(p-chlorophenyl)phosphine, tris(p-methoxyphenyl)phosphine, bis-(diphenylphosphino)methane, 1,2-bis-(diphenylphosphino)ethane, bis-(dicyclohexylphosphino)ethane, bis-(2-diphenylphosphinoethyl)phenylphosphine, 1,1,1-tris-(diphenylphosphinomethyl)ethane, tris-(2-diphenylphosphinoethyl)phosphine, 1,1,4,7,10,10-hexaphenyl-1,4,7,10-tetraphosphadecane, and 1,1-bis-(diphenylphosphino)ferrocene. The complex catalysts are used in conjunction with a ligand having a trivalent donor atom. The preferred ligand is phosphine or a phosphine derivative such as those listed above. The relative amounts of phosphine and Group VIII metal which can be used in the processes according to the invention are best expressed as a ratio of the number of moles of phosphorus in the phosphine compound to the number of moles of Group VIII metal. The preferred catalysts are salts, carbonyl compounds, chelates, or complexes of Rh, Pd, Ir, and Ru. Particularly preferred catalysts are RhCl_3 , PdCl_2 , PdBr_2 , IrCl_3 , $\text{Pd}(\text{OOCCH}_3)_2$, $(\text{RhCl}(\text{CO})_2)_2$, palladium bis triphenylphosphine, and palladium tris triphenyl phosphine. A more preferred Pd II catalyst is $\text{Pd}(\text{OAc})_2$.

The term "polyprotection" as used herein means reacting one or more protecting reagents at one or more heteroatoms.


Synthetic Schemes

Scheme 1 illustrates the general procedure for the preparation of the compounds of Formula 1 where each Y and each Z are independently selected from halogen, -OH, -OR, -O-C=O-R, n is equal to 0, 1, 2, 3, 4 or 5, m is equal to 0, 1, 2, 3, 4 or 5, each A and each B are independently selected from the group comprising alkyl with at least one carbon atom, aryl, aralkyl, P_n (wherein P_n is a protecting

group), and H. It should be appreciated that with regard to compounds of Formulas 1B, 1C, 1D, 1E, and 1F, A and B do not represent H; and, with regard to compounds of Formula 1A, when any Y or any Z is halogen, A or B is absent. Preferably, the sum of n and m is 1 or more. In some embodiments R is alkyl with at least two

5 carbon atoms.

Scheme 1.



Protection/Acylation of Benzoic Acid Halide Coupling Partner

10 As depicted in Scheme 1, the benzoic acid derivative 1A is optionally acylated and/or optionally protected with a desired number and type of acyl group(s) equal to or less than n and/or with the desired number and type of protecting group(s) equal to or less than n to yield benzoic acid derivative 1B where A is selected from the group comprising alkyl with at least one carbon atom, aryl,

15 aralkyl, and P_n .

A variety of reaction pathways starting with benzoic acid derivative 1A can be used to obtain the benzoic acid derivative 1B. These include homogenous peracylation or polyacylation (condition I), homogenous polyprotection (condition II), selective acylation followed by selective protection (condition III), selective

20 protection followed by selective acylation (condition IV), heterogeneous peracylation (condition V), heterogeneous polyprotection (condition VI), selective acylation followed by homogeneous or heterogeneous polyprotection (condition VII) and selective protection followed by homogeneous or heterogeneous

peracylation (condition IIX). It is contemplated that under condition VII, selective acylation may occur at one or more substituent sites and polyprotection may occur at one or more substituents sites. In like manner, it is contemplated that under condition IIX, selective protection may occur at one or more substituent sites and peracylation may occur at one or more substituents sites. When each Y or each Z is halogen, the acylation and/or protection and diprotection steps may be entirely omitted. Where one or more of Y and Z are halogen and one or more of Y and Z is neither H nor halogen, the acylation and/or protection and diprotection steps may be implemented.

10 It is to be understood from this description that benzoic acid 1A includes benzoic acid and monohydroxyl benzoic acids 2-hydroxy benzoic acid (salicylic acid), 3-hydroxy benzoic acid and 4-hydroxy benzoic acid in which case only conditions I or II can apply.


15 Benzoic acid 1A also includes but is not limited to the dihydroxyl benzoic acids. Illustrative examples of which include but are not limited to 2,3-dihydroxybenzoic acid, 3,5-dihydroxybenzoic acid (α -resorcylic acid), 2,4-dihydroxybenzoic acid (β -resorcylic acid), 2,6-dihydroxybenzoic acid (γ -resorcylic acid), 2,5-dihydroxybenzoic acid (hydroquinonecarboxylic acid, gentisic acid) and 3,4-dihydroxybenzoic acid (protocatechuic acid) in which case conditions I-VI can apply.

20 Benzoic acid 1A also includes but is not limited to the trihydroxyl benzoic acids. Illustrative examples of which include but are not limited to 3,4,5-trihydroxybenzoic acid (gallic acid), 2,3,4-trihydroxybenzoic acid, 2,4,5-trihydroxybenzoic acid and 2,4,6-trihydroxybenzoic acid (phloroglucinolcarboxylic acid) in which case conditions I-IIX can apply.

25 By way of illustration, the resorcylic acid analogs of 1A, where $n=2$ and each Y is OH, has two hydroxyls that can be 1) both esterified with one or more esterifying agents (condition I), 2) both protected with a one or more protecting agents (condition II), 3) selectively esterified with an esterifying agent and 30 selectively protected with a protecting agent (condition III), 4) selectively protected with a protecting agent and selectively esterified with an esterifying agent (condition IV), 5) heterogeneous diesterification with both hydroxyls esterified with two


different esterifying agents (condition V), or 6) heterogeneous diprotection with both hydroxyls protected with two different protecting agents (condition VI).

In the example of diesterification (condition I), both hydroxyls are reacted with at least 2 equivalents of an esterifying agent in a basic solvent (or in an inert solvent with added base) to give the diester resorcylic acid derivative 1B_a where A₁ and A₂ are the same. Preferably, a 5 molar equivalent excess of the esterifying agent is used.



In the example of diprotection (condition II), both hydroxyls are reacted with at least 2 equivalents of a protection agent and under conditions known to one skilled in the art for the desired alcohol protecting group to give diprotected resorcylic acid derivative 1B_a where A₁ and A₂ are the same.

In the example of selective esterification followed by selective protection (condition III), resorcylic acid is first reacted with 1 equivalent or more of an esterifying agent in a basic solvent (or in an inert solvent with added base) to give resorcylic acid derivative 1B_b where OA₁ represents an ester. Intermediate 1B_b is then reacted with 1 equivalent or more of a protection agent and under conditions known to one skilled in the art for the desired alcohol protecting group to give resorcylic acid derivative 1B_a where OA₂ represents a protected alcohol.



In the example of selective protection followed by selective esterification (condition IV), resorcylic acid is first reacted with 1 equivalent or more of a protection agent and under conditions known to one skilled in the art for the desired alcohol protecting group to give resorcylic acid derivative 1B_b where OA₁ represents a protected alcohol. Intermediate 1B_b is then reacted with 1 equivalent or more of an esterifying agent in a basic solvent (or in an inert solvent with added base) to give resorcylic acid derivative 1B_a where OA₂ represents an ester.

In the example of heterogeneous diesterification (condition V), resorcylic acid is first reacted with 1 equivalent or more of a first esterifying agent in a basic solvent (or in an inert solvent with added base) to give resorcylic acid derivative 1B_b where OA₁ represents a first ester. Intermediate 1B_b is then reacted with 1
5 equivalent or more of a second esterifying agent to give resorcylic acid derivative 1B_a where OA₂ represents a second ester.

In the example of heterogeneous diprotection (condition VI), resorcylic acid is first reacted with 1 equivalent or more of a first protection agent under conditions known to one skilled in the art for the desired alcohol protecting group to give
10 resorcylic acid derivative 1B_b where OA₁ represents a first protected alcohol. Intermediate 1B_b is then reacted with 1 equivalent or more of a second protecting agent to give resorcylic acid derivative 1B_a where OA₂ represents a second protected alcohol.

Suitable acylating agents useful in the step or steps described above include
15 but are not limited to acid anhydrides, acid halides, and mixed acid anhydrides of the formulas R₄CO₂COR₄, R₄COX, R₄CO₂COR₅ respectively where R₄ is alkyl with at least one carbon atom, aryl, aralkyl and diastereoisomers of the foregoing, R₅ is alkyl with at least one carbon atom, aryl, aralkyl and diastereoisomers of the foregoing, X is halogen. Cl is a preferred halogen. Acylation is carried out with
20 basic solvents including but not limited to ethyl amine, diethyl amine, triethylamine, pyrrole, imidazole, triazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, pyrazine, triazine, pyrrolidine, piperidine, DMF, NMP, dimethylpropyl urea, hexamethyl phosphotriamide (HMPA), and mixtures of said solvents. Other appropriate solvents are known to one skilled in the art. Pyridine is a preferred basic
25 solvent. Alternatively, acylation can be carried out in an inert solvent with added base.

In the examples where acylation results in an ester (i.e. X is O), esterification is followed by treatment with aqueous formic acid and optionally by recrystallization. In the examples where acylation results in an amide (i.e. X is NH),
30 acylation is followed by treatment with any mild acid and optionally by recrystallization.

Suitable protecting groups are described in T. W. Greene, Protective Groups in Organic Synthesis, John Wiley & Sons, New York, N.Y., 1991, 2nd Edition.

Preferred alcohol protecting groups are those that can be selectively cleaved over a phenyl ester and include those derived from chloromethoxymethyl acetate (MOMCl), chloroacetoxy anhydride, levulonic anhydride. Silicon protecting groups such as *tert*-butyldimethylsilyl and *tert*-butylsilyl were not found to be compatible.

5 Ether protecting groups such as methyl and benzyl ethers do not work as well since the Lewis acid conditions employed during deprotection are harsher as to other portions of the desired products.

Preferred amine protecting groups are those that can be selectively cleaved over a phenyl amine and include *tert*-butyloxycarbonyl (BOC),

10 fluorinomethyloxycarbonyl (Fmoc), and benzyloxycarbonyl (Cbz).

Halogenation of Acid Chloride Coupling Partner

Returning to Scheme 1, benzoic acid derivative 1B is reacted with a halogenating agent in an inert solvent to yield the benzoic acid halide derivative 1C where X is any halogen.

15 Suitable halogenating agents useful in this step include but are not limited to any of the halogenating agents known to the art, such as thionyl halides including thionyl chloride, thionyl bromide and thionyl iodide, oxalyl halides including oxalyl chloride, oxalyl bromide and oxalyl iodide, phosphorous tri- and pentahalides and phosphorous oxyhalides. Halogenation is carried out in an inert solvent including

20 but not limited to halogenated alkanes such as methylene chloride and chloroform; cyclic ethers, such as THF and dioxane; N,N-dialkylsubstituted acylamides, such as DMF and DMAC; and mixtures of said solvents.

The halogenation reaction may optionally include a buffering agent capable of associating with HX acid. Suitable buffering agents include weak bases.

25 Benzotriazole is a preferred buffering agent during halogenation because it precipitates as a salt after reacting with a HX acid.

The reaction is effected at a temperature from about 0° to about 100° C. Preferably, the reaction is effected using thionyl chloride or oxalyl chloride in a polar solvent such as methylene chloride, at 80°C, in the presence of benzotriazole

30 to yield the corresponding acid chloride.

Optionally, the intermediate 1C is recrystallized from hexane.

Preparation of Styrene Coupling Partner

Also depicted in Scheme 1 is styrene derivative 1D. Styrene 1D is optionally acylated and/or optionally protected with a desired number and type of acyl group(s) equal to or less than m and/or with the desired number and type of protecting
5 group(s) equal to or less than m to yield styrene benzoic acid derivative 1E where A is selected from the group comprising alkyl with at least one carbon atom, aryl, aralkyl, and P_m.

A variety of reaction pathways starting with styrene derivative 1D can be used to obtain the styrene derivative 1E. These pathways include homogenous
10 peracylation or polyacylation (condition I), homogenous polyprotection (condition II), selective acylation followed by selective protection (condition III), selective protection followed by selective acylation (condition IV), heterogeneous peracylation (condition V), heterogeneous polyprotection (condition VI), selective acylation followed by homogeneous or heterogeneous polyprotection (condition
15 VII) and selective protection followed by homogeneous or heterogeneous peracylation (condition IIX). It is contemplated that under condition VII, selective acylation may occur at one or more substituent sites and polyprotection may occur at one or more substituents sites. In like manner, it is contemplated that under condition IIX, selective protection may occur at one or more substituent sites and
20 peracylation may occur at one or more substituents sites.

It is to be understood from this description that styrene derivative 1A includes styrene and the monohydroxyl styrenes 2-hydroxy styrene, 3-hydroxy styrene and 4-hydroxy styrene in which case only conditions I or II can apply. Styrene derivative 1D also includes but is not limited to the dihydroxyl styrenes.
25 Illustrative examples of which include but are not limited to 2,3-dihydroxystyrene, 3,5-dihydroxystyrene, 2,4-dihydroxystyrene, 2,6-dihydroxystyrene, 2,5-dihydroxystyrene and 3,4-dihydroxystyrene in which case conditions I-VI can apply. Styrene derivative 1D also includes but is not limited to the trihydroxyl styrenes. Illustrative examples of which include but are not limited to 3,4,5-trihydroxystyrene,
30 2,3,4-trihydroxystyrene, 2,4,5-trihydroxystyrene and 2,4,6-trihydroxystyrene in which case conditions I-IIX can apply.

By way of illustration, the monohydroxyl styrene analogs of 1D (where n=1 and Y is OH) has one hydroxyl that can be 1) esterified with an esterifying agent

(condition I) or 2) protected with a protecting agent (condition II). The monohydroxyl styrenes can be prepared by saponification of the corresponding acetoxy styrenes, e.g. 4-hydroxystyrene from 4-acetoxystyrene.

Coupling Step via Decarbonylative Heck Reaction

5 Returning to Scheme 1, benzoic acid halide 1C is reacted with styrene 1E via a decarbonylative coupling reaction to give stilbene 1F. The reaction conditions associated with the coupling reaction can include an inert apolar, high-boiling point aromatic or hydrocarbon solvent with a transition metal catalyst, N-heterocyclic carbene-type ligand and base.

10 Suitable inert solvents useful in the coupling step include but are not limited to benzene, toluene, xylene, chlorobenzene, decane, dodecane, and mixtures of those solvents. Xylene is a preferred solvent in the coupling step.

A preferred transition metal catalyst used in the coupling step is $\text{Pd}(\text{OAc})_2$.

15 A preferred N-heterocyclic carbene-type ligands useful in the coupling step is N,N-bis-(2,6-diisopropylphenyl) 4,5-dihydroimidazolium chloride in an amount of 0.1-10 mol %, preferably 5-10 mol%.

Surprisingly, the nature of the added base significantly affects the efficiency of the reaction. Non-coordinating amine bases result in greater yields than coordinating amine bases. Examples of non-coordinating bases useful in the process include but are not limited to N-ethylmorpholine (NEM), N,N-dimethylbenzylamine, dimethylaminopyridine (DMAP), proton sponge (1,8-diaminonaphthalene), diisopropylethylamine (Hunig's base) and triethylamine. N-ethylmorpholine and N,N-dimethylbenzylamine are preferred bases in the coupling step. Use of smaller amines capable of transition metal coordination still work to
25 obtain the coupled product 1F but are less efficient. Furthermore, phosphine ligands can also be added to the reaction; however, added phosphine ligand inhibits the reaction giving greatly lowered yields.

The reaction is effected at a temperature from about 0 to about 150° C. Preferably, the reaction is refluxed in xylene.

30 The reaction depicted in Scheme 1 is concluded using standard work-up conditions and the reaction products can be purified using a variety of techniques known to those skilled in the art including silica gel chromatography.

Deprotection Step

After performing the coupling reaction, in those cases where one or more protecting groups exist on the stilbene derivative 1F, the protection groups may be deprotected as described in T. W. Greene, Protective Groups in Organic Synthesis, John Wiley & Sons, New York, N.Y., 1991, 2nd Edition for each of the respective protection groups and may require one or more steps to complete. With respect to the preferred hydroxyl protecting groups, the methoxymethyl acetate (MOM) may be selectively hydrolyzed by reaction with sodium iodide and trimethylsilyl chloride (TMSCl); chloroacetoxy groups may be selectively hydrolyzed when treated with 50% aqueous pyridine at a pH of 6.7; the levulinoate (lev) may be selectively hydrolyzed with sodium thiosulfate (Na₂SO₃) and sodium metabisulfite.

If desired, phenolic esters may also be cleaved, if present, by hydrolysis using saponification techniques known to one skilled in the art. Ester hydrolysis may be carried out before or after deprotection of any protecting groups.

In the example of stilbene derivatives 1F akin to resveratrol, where Y=O at the 3 and 5 positions, n=2, Z=O at the 4' position and m=1, the derivative may be 1) deesterified at all three phenolic sites (3,5, and 4'), 2) deprotected and/or deesterified at all three phenolic sites and 3) selectively deprotected at any of the three phenolic sites.


In the example of deesterification at any of the three phenolic sites, basic hydrolysis (saponification) can be carried out in a polar aprotic inert solvent including but not limited to dimethylsulfoxide (DMSO), N-methylpyrrolidone, tetrahydrofuran (THF), acetone, acetonitrile, methylene chloride, chloroform, mixtures of said polar aprotic inert solvents and others known to one skilled in the art with a suitable base including but not limited to alkali metal hydroxides (e.g., LiOH, NaOH and KOH) and alkaline earth metal hydroxides (e.g., Mg(OH)₂, Ca(OH)₂, and Ba(OH)₂). A preferred solvent is THF. A preferred alkaline earth metal hydroxide base is KOH.

Olefin Isomerization

The stilbene compounds of the present invention possess a conjugated double bond. The double bond may exist in the *cis* or *trans* configuration. The present invention includes both *cis* and *trans* isomers as mixtures or in purified form. Furthermore, one aspect of the present invention includes a process for

isomerizing the *trans* configuration to the *cis* configuration. As depicted in Scheme 2, the isomerization step may be conducted to the stilbene derivatives when Y and/or Z are acylated, protected, or the free alcohol. Compounds of the Formulas 1A3, 1B, I, II, 1, 3-7 can be diluted in an inert solvent and irradiated with ultraviolet light, preferably of wavelength $\lambda = 280-350$ nm. This ultraviolet irradiation is carried in organic solution, at temperatures of 0° to 50° , preferably of 10° to 30° . The irradiation can be carried out with the aid of the most diverse commercial ultraviolet apparatuses. By way of example, irradiation can be carried out with a medium-pressure mercury vapor lamp through a pyrex filter. The irradiation time is generally from 2 to 100 hours, preferably 5 to 50 hours, especially 10 to 20 hours.

Scheme 2.



Resveratrol Esters Coupling Step via Decarbonylative Heck reaction

In the preferred embodiments that follow, selective ester derivatives of resveratrol can be prepared where the 3-hydroxyl, 5-hydroxyl and 4'-hydroxyl can independently be selectively substituted. In the preferred ester derivatives of resveratrol, R_1 , R_2 and R_3 are each independently selected from the group comprising alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing and P_1 , P_2 and P_3 are alcohol protecting groups. In some embodiments R is alkyl with at least two carbon atoms. In some embodiments the sum of m and n is greater than or equal to one but less than nine. In other embodiments the sum of m and n is three or more but less than nine.


With reference to Formula 1A₃, a first preferred embodiment constitutes the 3-ester derivatives of resveratrol. In that embodiment, A_1 is R_1 , A_2 is P_2 or OH, and A_3 is P_3 or OH. As explained more specifically later in this description, this embodiment is equivalent (or degenerate) when A_1 is P_1 or OH, A_2 is R_2 , and A_3 is P_3 or OH. A second preferred embodiment constitutes the 3,5-diester derivatives of resveratrol. Therein, A_1 is R_1 , A_2 is R_2 , and A_3 is P_3 or OH. A third preferred embodiment constitutes the 3,4'-diester derivatives of resveratrol when A_1 is R_1 , A_2 is P_2 or OH, and A_3 is R_3 . As explained more specifically later in this description,

this embodiment is equivalent (or degenerate) when A_1 is P_1 or OH, A_2 is R_2 , and A_3 is R_3 . A fourth preferred embodiment constitutes the 4'-ester derivatives of resveratrol. Therein, A_1 is P_1 or OH, A_2 is P_2 or OH, and A_3 is R_3 . A fifth preferred embodiment constitutes the 3,5,4'-triester derivatives of resveratrol. In the

5 esters may be the same or different. Furthermore, in the embodiments that follow with more than one protecting group, P_1 , P_2 and P_3 may be the same or different. The resveratrol ester derivatives can be of the *cis*-stilbene or *trans*-stilbene configuration.

10 Scheme 3 illustrates one procedure for the preparation of the mono- and di-esters of resveratrol.


Scheme 3.



Selective 3-esters of Resveratrol

15 Selective 3-ester derivatives of resveratrol may also be prepared by the process depicted in Scheme 4.

Scheme 4.



In a preferred embodiment, 3-ester compounds of Formula 3, pharmaceutically and cosmetically acceptable salts thereof wherein R₁ is selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing, are synthesized from the steps a and b (or alternatively c and d), e, f, and g: wherein step a) is esterifying resorcylic acid with an esterifying agent in solution or in suspension of a first solvent to give the 3-ester of resorcylic acid of Formula 3A; step b) is reacting the hydroxyl of said compound of Formula 3A with a first alcohol protecting agent in solution or in suspension of a second solvent to give a compound of Formula 3B; where P₁ is a first alcohol protecting group; step c) is reacting resorcylic acid with a first alcohol protecting agent in solution or in suspension of a second solvent to give a compound of Formula 3C, where P₁ is a first alcohol protecting group; step d) is reacting the 5-hydroxy of said compound of Formula 3C with an esterifying agent in solution or in suspension of a first solvent to give the 3-ester of resorcylic acid of Formula 3B; step e) is halogenating said compound of Formula 3B using a halogenating agent in solution or in suspension of a third solvent to give the acid halide of Formula 3D, wherein X is halogen; step f) is coupling said compound of Formula 3B with a compound of Formula 3D, where P₂

is a second alcohol protecting group, with Pd II catalyst, N-heterocyclic carbene-type ligand, and a first base in solution or in suspension of a fourth solvent to yield a compound of Formula 3E; and, step g) is deprotecting said first and second alcohol protecting groups from said compound of Formula 3E to give a compound of Formula 3. In some embodiments R is alkyl with at least two carbon atoms.

Compounds of the invention are those prepared by the above process.

As mentioned previously, Scheme 4 illustrates the general procedure for the preparation of the compounds of Formula 3. The compounds of Formula 3 may be generally characterized as the 3-esters of resveratrol, that is, the phenolic hydroxyls at the 5 and 4' positions are the free alcohols while the phenolic hydroxyl at the 3 position is an ester.


The 3-esters of resveratrol are prepared by either 1) reacting one of the resorcylic acid hydroxyls with one or more equivalents of a desired esterifying agent followed by protection of the remaining hydroxyl with a suitable alcohol protection agent or 2) reacting one of the resorcylic acid hydroxyls with one or more equivalents of a suitable protecting agent followed by esterification of the remaining hydroxyl with one or more equivalents of a desired esterifying agent. The esterified/protected acid is halogenated to a corresponding acid halide. A 4-protectedhydroxy styrene is made by reacting 4-hydroxystyrene with a desired protecting agent. The 4-protectedhydroxy styrene is reacted with the esterified/protected acid halide in a decarbonylative Heck reaction as described above. After coupling, the protection groups at the 5 and 4' positions are removed to give the 3-ester of resveratrol.

In either of the above reaction pathways, reaction at one of the two phenolic hydroxyls of resorcylic acid is equivalent (or degenerate) to reaction at the other hydroxyl because of the symmetry of α -resorcylic acid. By way of example, if the first reaction is monoesterification of α -resorcylic acid, reaction at the 3-hydroxyl would be equivalent to the reaction at the 5-hydroxyl as both products would be indistinguishable because of the mirror plane passing along the first and fourth carbons of the aromatic ring. In that regard, compound 3-acetoxy-5-hydroxy benzoic acid is equivalent to the compound 5-acetoxy-3-hydroxy benzoic acid. Therefore, the 5-ester derivatives of resveratrol are equivalent (or degenerate) to the 3-ester derivatives of resveratrol.

Selective 3,5-diester of Resveratrol

Selective 3,5-diester derivatives of resveratrol may also be prepared by the process depicted in Scheme 5.

Scheme 5.



5

In a preferred embodiment, 3,5-diester compounds of Formula 4, pharmaceutically and cosmetically acceptable salts thereof wherein R_1 and R_2 are independently selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing, are synthesized from the steps of a (or alternatively b and c), d, e, and f: wherein step a) is esterifying resorcylic acid with an esterifying agent in solution or in suspension of a first solvent to give the 3,5-diester of resorcylic acid of Formula 4A wherein R_1 and R_2 are the same; step b) is esterifying resorcylic acid with a first esterifying agent in solution or in suspension of a first solvent to give the 3-ester of Formula 4B; step c) is esterifying said compound of Formula 4B with a second esterifying agent in solution or in suspension of a second solvent to give the 3,5-diester of Formula 4A wherein R_1 and R_2 are different; step d) is halogenating said compound of Formula 4A using a halogenating agent in solution or in suspension of a third solvent to give the acid halide of Formula 4C, wherein X is halogen; step e) is coupling said compound of

Formula 4B with a compound of Formula 4D, wherein P₁ is a first alcohol protecting group, with Pd II catalyst, N-heterocyclic carbene-type ligand, and a first base in solution or in suspension of a fourth solvent to yield a compound of Formula 4E; and, step f) is deprotecting said first alcohol protecting group from said
5 compound of Formula 4E to give a compound of Formula 4. In some embodiments R is alkyl with at least two carbon atoms.

Compounds of the invention are those prepared by the above process.


Returning to Scheme 5, the general procedure for the preparation of the compounds of Formula 4 is depicted. The compounds of Formula 4 may be
10 generally characterized as the 3,5-diesters of resveratrol, that is, the phenolic hydroxyl at the 4' position is the free alcohol while the phenolic hydroxyls at the 3 and 5 positions are esters. In some embodiments the 3 and 5 esters are the same while in other embodiments the 3 and 5 esters are different.

The 3,5-diesters of resveratrol are prepared by reacting resorcylic acid with
15 two or more equivalents of a desired esterifying agent. In the alternative for mixed esters, resorcylic acid is reacted with one or more equivalents of a first esterifying agent and subsequently reacted with one or more equivalents of a second esterifying agent. Advantageously, the mixed ester can be prepared in one pot if the second esterifying agent is added after the first esterifying agent has reacted to completion
20 or near to completion. The diesterified (heterogeneous or homogenous esters) resorcylic acid derivative is halogenated to a corresponding acid halide. A 4-protectedhydroxy styrene is made by reacting 4-hydroxystyrene with a desired protecting agent. The 4-protectedhydroxy styrene is reacted with the diesterified acid chloride in a decarbonylative Heck reaction as described above. After
25 coupling, the protection group at the 4' position is removed to give the 3,5-diesters of resveratrol. A preferred protection group used with the 4-hydroxy styrene is chloroacetoxy.

Selective 3,4'-diesters of Resveratrol

Selective 3,4'-diester derivatives of resveratrol may also be prepared by the
30 process depicted in Scheme 6.

Scheme 6.



In another preferred embodiment, 3,4'-diester compounds of Formula 5, pharmaceutically and cosmetically acceptable salts thereof wherein R_1 and R_3 are independently selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing, are synthesized from the steps comprising the steps a and b (or alternatively c and d), and e, f, and g: wherein step a) is esterifying resorcylic acid with an esterifying agent in solution or in suspension of a first solvent to give the 3-ester of resorcylic acid of Formula 5A; step b) is reacting the 5-hydroxy of said compound of Formula 5A with a first alcohol protecting agent in solution or in suspension of a second solvent to give a compound of Formula 5B, where P_2 is a first alcohol protecting group; step c) is reacting resorcylic acid with a first alcohol protecting agent in solution or in suspension of a second solvent to give a compound of Formula 5C, where P_2 is an alcohol protecting group; step d) is reacting the 5-hydroxy of said compound of Formula 5C with an esterifying agent in solution or in suspension of a first solvent to give the 3-ester of resorcylic acid of Formula 5B; step e) is halogenating said compound of Formula 5B using a halogenating agent in solution or in suspension of a third solvent to give the acid halide of Formula 5D, wherein X is halogen; step f) is coupling said compound of Formula 5D with a compound of Formula 5E, with Pd II catalyst, N-heterocyclic

carbene-type ligand, and a first base in solution or in suspension of a fourth solvent to yield a compound of Formula 5E; and, step g) is deprotecting said first and second protecting groups from said compound of Formula 5E to give a compound of Formula 5. In some embodiments R is alkyl with at least two carbon atoms.

5 Compounds of the invention are those prepared by the above process.

Returning to Scheme 6, the general procedure for the preparation of the compounds of Formula 5 is shown. The compounds of Formula 5 may be generally characterized as the 3,4'-diesters of resveratrol, that is, the phenolic hydroxyl at the 5 position is the free alcohol while the phenolic hydroxyls at the 3 and 4' positions
10 are esters.


The 3,4'-diesters of resveratrol are prepared by either 1) reacting one of the resorcylic acid hydroxyls with one or more equivalents of a desired esterifying agent followed by protection of the remaining hydroxyl with a suitable alcohol protection agent or 2) reacting one of the resorcylic acid hydroxyls with one or more
15 equivalents of a suitable protecting agent followed by esterification of the remaining hydroxyl with one or more equivalents of a desired esterifying agent. The esterified/protected acid is halogenated to a corresponding acid halide. A 4-ester styrene is made by reacting 4-hydroxystyrene with a desired esterifying agent. The 4-ester styrene is reacted with the esterified/protected acid halide in a
20 decarbonylative Heck reaction as described above. After coupling, the protection group at the 5 position is removed to give the 3,4'-diester of resveratrol.

As mentioned in relation to the synthesis of the 3-ester (and degenerate 5-ester) derivatives of resveratrol earlier, in either of the above reaction pathways, reaction at one of the two phenolic hydroxyls of resorcylic acid is equivalent (or
25 degenerate) to reaction at the other hydroxyl because of the symmetry of α -resorcylic acid. Hence, the 5,4'-diester derivatives of resveratrol are equivalent (or degenerate) to the 3,4'-diester derivatives of resveratrol.

Selective 4'-esters of Resveratrol

Selective 4'-ester derivatives of resveratrol may also be prepared by the
30 process depicted in Scheme 7.

Scheme 7.



- In another preferred embodiment, 4'-ester compounds of Formula 6, pharmaceutically and cosmetically acceptable salts thereof wherein R₃ is selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing, are synthesized from the steps of a, b (or in the alternative c and d), e, f, and g: wherein step a) is esterifying 4-hydroxy styrene with an esterifying agent in solution or in suspension of a first solvent to give the 4-ester styrene of Formula 6D; step b) is reacting resorcylic acid with an alcohol protecting agent in solution or in suspension of a second solvent to give a compound of Formula 6B; wherein P₁ is a first alcohol protecting group and P₂ is a second alcohol protecting group; step c) reacting resorcylic acid with a first alcohol protecting agent in solution or in suspension of a second solvent to give a compound of Formula 6A; step d) is reacting a compound of Formula 6A with a second alcohol protecting agent in solution or in suspension of a third solvent to give a compound of Formula 6B where P₁ and P₂ are different; step e) is halogenating said compound of Formula 6B using a halogenating agent in solution or in suspension of a fourth solvent to give the acid halide of Formula 6C, wherein X is halogen; step f) is coupling said compound of Formula 6D with a compound of Formula 6C with Pd II catalyst, N-heterocyclic carbene-type ligand, and a first base in solution or in suspension of a fifth solvent to yield a compound of Formula 6E; and, step e) is deprotecting said first alcohol protecting group from said compound of Formula 6D to give a

compound of Formula 6. In some embodiments R is alkyl with at least two carbon atoms.

Compounds of the invention are those prepared by the above process.


Returning to Scheme 7, the general procedure for the preparation of the compounds of Formula 6 is shown. The compounds of Formula 6 may be generally characterized as the 4'-esters of resveratrol, that is, the phenolic hydroxyls at the 3 and 5 positions are free alcohols while the phenolic hydroxyl at the 4' position is an ester.

The 4'-esters of resveratrol are prepared by reacting resorcylic acid with two or more equivalents of a suitable protecting agent. The diprotected acid is halogenated to a corresponding acid halide. A 4-ester styrene is made by reacting 4-hydroxystyrene with a desired esterifying agent. The 4-ester styrene is reacted with the protected acid halide in a decarbonylative Heck reaction as described above. After coupling, the protection groups at the 3 and 5 positions are removed to give the 4'-ester of resveratrol.

Selective 3,5,4'-triesters of Resveratrol

Selective 3,5,4'-triester derivatives of resveratrol may also be prepared by the process depicted in Scheme 8.

Scheme 8.



20

In another preferred embodiment, 3,5,4'-triester compounds of Formula 7, pharmaceutically and cosmetically acceptable salts thereof wherein R_1 , R_2 and R_3

are independently selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing, are synthesized from the steps of a (or alternatively b and c), d, e, f, and g: wherein step a) is esterifying resorcylic acid with an esterifying agent in solution or in suspension of a first solvent to give the
5 3,5-diester of resorcylic acid of Formula 7A; step b) is esterifying resorcylic acid with a first esterifying agent in solution or in suspension of a first solvent to give the 3-ester of Formula 7B step c) is esterifying said compound of Formula 7B with a second esterifying agent in solution or in suspension of a second solvent to give the 3,5-diester of Formula 7A where R_1 and R_2 are different; step d) is halogenating said
10 compound of Formula 7A using a halogenating agent in solution or in suspension of a third solvent to give the acid halide of Formula 7C, wherein X is halogen; step e) is coupling said compound of Formula 7D with a compound of Formula 7C with Pd II catalyst, N-heterocyclic carbene-type ligand, and a first base in solution or in suspension of a fourth solvent to yield said compound of Formula 7. In some
15 embodiments R is alkyl with at least two carbon atoms.

Compounds of the invention are those prepared by the above process.

Returning to Scheme 8, the general procedure for the preparation of the compounds of Formula 7 is shown. The compounds of Formula 7 may be generally characterized as the 3,5,4'-triesters of resveratrol, that is, the phenolic hydroxyls at
20 the 3,5,4' positions are esters.

The 3,5,4'-triesters of resveratrol are prepared by reacting resorcylic acid with two or more equivalents of a desired esterifying agent. The diesterified resorcylic acid is halogenated to a corresponding acid halide. A 4-ester styrene is made by reacting 4-hydroxystyrene with a desired esterifying agent. The 4-ester
25 styrene is reacted with the diesterified acid halide in a decarbonylative Heck reaction as described above to give the 3,5,4'-triester of resveratrol.

Preparation of Resveratrol Esters

The processes described to produce the 3-esters, 3,5-diesters, 3,4'-diesters, 4'-esters, and 3,5,4'-triesters of resveratrol can be used to produce resveratrol itself
30 by augmenting the respective processes with an additional step: hydrolysis with an alkaline earth hydroxide or alkaline earth alkoxide in a polar aprotic solvent followed by acidification.

Preferred Compounds

Compounds of the invention are those prepared by the processes above, especially those selected from: 5,4'-dihydroxy-3-acetoxy stilbene (3-acetoxy resveratrol), 3,5-dihydroxy-4'-acetoxy stilbene (4'-acetoxy resveratrol), 3,4'-

5 dihydroxy-5-acetoxy stilbene (5-acetoxy resveratrol), 4'-hydroxy-3,5-diacetoxy stilbene (3,5-diacetoxy resveratrol), 5-hydroxy-3,4'-diacetoxy stilbene (3,4'-diacetoxy resveratrol), 3,5,4'-triaceoxy stilbene (3,5,4'-triaceoxy resveratrol); 5,4'-dihydroxy-3-propanoate stilbene, 3,5-dihydroxy-4'-propanoate stilbene, 3,4'-dihydroxy-5-propanoate stilbene, 4'-hydroxy-3,5-dipropanoate stilbene, 5-hydroxy-

10 3,4'-dipropanoate stilbene, 3,5,4'-tripropanoate stilbene; 5,4'-dihydroxy-3-butanoate stilbene, 3,5-dihydroxy-4'-butanoate stilbene, 3,4'-dihydroxy-5-butanoate stilbene, 4'-hydroxy-3,5-dibutanoate stilbene, 5-hydroxy-3,4'-dibutanoate stilbene, 3,5,4'-tributanoate stilbene; 5,4'-dihydroxy-3-pentanoate stilbene, 3,5-dihydroxy-4'-pentanoate stilbene, 3,4'-dihydroxy-5-pentanoate stilbene, 4'-hydroxy-3,5-

15 dipentanoate stilbene, 5-hydroxy-3,4'-dipentanoate stilbene, 3,5,4'-tripentanoate stilbene; 5,4'-dihydroxy-3-hexanoate stilbene, 3,5-dihydroxy-4'-hexanoate stilbene, 3,4'-dihydroxy-5-hexanoate stilbene, 4'-hydroxy-3,5-dihexanoate stilbene, 5-hydroxy-3,4'-dihexanoate stilbene, 3,5,4'-trihexanoate stilbene; sorbic fatty acid derivatives including but not limited to: 5,4'-dihydroxy-3-(2,4-hexadienoate)

20 stilbene, 3,5-dihydroxy-4'-(2,4-hexadienoate) stilbene, 3,4'-dihydroxy-5-(2,4-hexadienoate) stilbene, 4'-hydroxy-3,5-(2,4-hexadienoate) stilbene, 5-hydroxy-3,4'-(2,4-hexadienoate) stilbene, 3,5,4'-tri-(2,4-hexanoate) stilbene; lauric fatty acid derivatives including but not limited to: 5,4'-dihydroxy-3-dodecanoate stilbene, 3,5-dihydroxy-4'-dodecanoate stilbene, 3,4'-dihydroxy-5-dodecanoate stilbene, 4'-

25 hydroxy-3,5-dodecanoate stilbene, 5-hydroxy-3,4'-dodecanoate stilbene, 3,5,4'-tridodecanoate stilbene; palmitic fatty acid derivatives including but not limited to: 5,4'-dihydroxy-3-hexadecanoate stilbene, 3,5-dihydroxy-4'-hexadecanoate stilbene, 3,4'-dihydroxy-5-hexadecanoate stilbene, 4'-hydroxy-3,5-hexadecanoate stilbene, 5-hydroxy-3,4'-hexadecanoate stilbene, 3,5,4'-trihexadecanoate stilbene; stearic fatty

30 acid derivatives including but not limited to: 5,4'-dihydroxy-3-octadecanoate stilbene, 3,5-dihydroxy-4'-octadecanoate stilbene, 3,4'-dihydroxy-5-octadecanoate stilbene, 4'-hydroxy-3,5-octadecanoate stilbene, 5-hydroxy-3,4'-octadecanoate stilbene, 3,5,4'-trioctadecanoate stilbene; oleic fatty acid derivatives including but

not limited to: 5,4'-dihydroxy-3-(9-octadecenoate) stilbene, 3,5-dihydroxy-4'-(9-octadecenoate) stilbene, 3,4'-dihydroxy-5-(9-octadecenoate) stilbene, 4'-hydroxy-3,5-(9-octadecenoate) stilbene, 5-hydroxy-3,4'-(9-octadecenoate) stilbene, 3,5,4'-tri-(9-octadecenoate) stilbene; linoleic fatty acid derivatives including but not limited to: 5,4'-dihydroxy-3-(9,12-octadecadienoate) stilbene, 3,5-dihydroxy-4'-(9,12-octadecadienoate) stilbene, 3,4'-dihydroxy-5-(9,12-octadecadienoate) stilbene, 4'-hydroxy-3,5-(9,12-octadecadienoate) stilbene, 5-hydroxy-3,4'-(9,12-octadecadienoate) stilbene, 3,5,4'-tri-(9,12-octadecadienoate) stilbene; linolenic fatty acid derivatives including but not limited to: 5,4'-dihydroxy-3-(6,9,12-octadecatrienoate) stilbene, 3,5-dihydroxy-4'-(6,9,12-octadecatrienoate) stilbene, 3,4'-dihydroxy-5-(6,9,12-octadecatrienoate) stilbene, 4'-hydroxy-3,5-(6,9,12-octadecatrienoate) stilbene, 5-hydroxy-3,4'-(6,9,12-octadecatrienoate) stilbene, 3,5,4'-tri-(6,9,12-octadecatrienoate) stilbene; linolenic fatty acid derivatives including but not limited to: 5,4'-dihydroxy-3-(9,12,15-octadecatrienoate) stilbene, 3,5-dihydroxy-4'-(9,12,15-octadecatrienoate) stilbene, 3,4'-dihydroxy-5-(9,12,15-octadecatrienoate) stilbene, 4'-hydroxy-3,5-(9,12,15-octadecatrienoate) stilbene, 5-hydroxy-3,4'-(9,12,15-octadecatrienoate) stilbene, 3,5,4'-tri-(9,12,15-octadecatrienoate) stilbene; α -linolenic fatty acid derivatives including but not limited to: 5,4'-dihydroxy-3-(3,6,9-octadecatrienoate) stilbene, 3,5-dihydroxy-4'-(3,6,9-octadecatrienoate) stilbene, 3,4'-dihydroxy-5-(3,6,9-octadecatrienoate) stilbene, 4'-hydroxy-3,5-dihydroxy-(3,6,9-octadecatrienoate) stilbene, 5-hydroxy-3,4'-(3,6,9-octadecatrienoate) stilbene, 3,5,4'-tri-(3,6,9-octadecatrienoate) stilbene; arachidonic fatty acid derivatives including but not limited to: 5,4'-dihydroxy-3-(5,8,11,14-eicosatetraenoate) stilbene, 3,5-dihydroxy-4'-(5,8,11,14-eicosatetraenoate) stilbene, 3,4'-dihydroxy-5-(5,8,11,14-eicosatetraenoate) stilbene, 4'-hydroxy-3,5-(5,8,11,14-eicosatetraenoate) stilbene, 5-hydroxy-3,4'-(5,8,11,14-eicosatetraenoate) stilbene, 3,5,4'-tri-(5,8,11,14-eicosatetraenoate) stilbene; eicosapentaenoic fatty acid derivatives including but not limited to: 5,4'-dihydroxy-3-(5,8,11,14,17-eicosapentaenoate) stilbene, 3,5-dihydroxy-4'-(5,8,11,14,17-eicosapentaenoate) stilbene, 3,4'-dihydroxy-5-(5,8,11,14,17-eicosapentaenoate) stilbene, 4'-hydroxy-3,5-(5,8,11,14,17-eicosapentaenoate) stilbene, 5-hydroxy-3,4'-(5,8,11,14,17-eicosapentaenoate) stilbene, 3,5,4'-tri-(5,8,11,14,17-eicosapentaenoate) stilbene; docosahexanoic fatty acid derivatives including but not

limited to: 5,4'-dihydroxy-3-(4,7,10,13,16,19-docosaehaenoate) stilbene, 3,5-dihydroxy-4'-(4,7,10,13,16,19-docosaehaenoate) stilbene, 3,4'-dihydroxy-5-(4,7,10,13,16,19-docosaehaenoate) stilbene, 4'-hydroxy-3,5-(4,7,10,13,16,19-docosaehaenoate) stilbene, 5-hydroxy-3,4'-(4,7,10,13,16,19-docosaehaenoate) stilbene, and 3,5,4'-tri-(4,7,10,13,16,19-docosaehaenoate) stilbene.


Compounds of the invention are also those esters of resorcylic acid halide useful for decarbonylative Heck reactions described above, especially those selected from: 3,5-diacetoxybenzoyl chloride, 3,5-diacetoxybenzoyl bromide, 3,5-diacetoxybenzoyl iodide, 3,5-dipropionate benzoyl chloride, 3,5-dipropionate benzoyl bromide, 3,5-dipropionate benzoyl iodide, 3,5-dibutanoate benzoyl chloride, 3,5-dibutanoate benzoyl bromide, 3,5-dibutanoate benzoyl iodide, C5: 3,5-dipentanoate benzoyl chloride, 3,5-dipentanoate benzoyl bromide, 3,5-dipentanoate benzoyl iodide, C6: 3,5-dihexanoate benzoyl chloride, 3,5-dihexanoate benzoyl bromide, 3,5-dihexanoate benzoyl iodide, sorbic: 3,5-dihexadienoate benzoyl chloride, 3,5-dihexadienoate benzoyl bromide, 3,5-dihexadienoate benzoyl iodide, lauric: 3,5-didodecanoate benzoyl chloride, 3,5-didodecanoate benzoyl bromide, 3,5-didodecanoate benzoyl iodide, palmitic: 3,5-dihexadecanoate benzoyl chloride, 3,5-dihexadecanoate benzoyl bromide, 3,5-dihexadecanoate benzoyl iodide, stearic: 3,5-dioctadecanoate benzoyl chloride, 3,5-dioctadecanoate benzoyl bromide, 3,5-dioctadecanoate benzoyl iodide, oleic: 3,5-di-(9-octadecenoate) benzoyl chloride, 3,5-di-(9-octadecenoate) benzoyl bromide, 3,5-di-(9-octadecenoate) benzoyl iodide, linoleic: 3,5-di-(9,12-octadecadienoate) benzoyl chloride, 3,5-di-(9,12-octadecadienoate) benzoyl bromide, 3,5-di-(9,12-octadecadienoate) benzoyl iodide, linolenic: 3,5-di-(6,9,12-octadecatrienoate) benzoyl chloride, 3,5-di-(6,9,12-octadecatrienoate) benzoyl bromide, 3,5-di-(6,9,12-octadecatrienoate) benzoyl iodide, 3,5-di-(9,12,15-octadecatrienoate) benzoyl chloride, 3,5-di-(9,12,15-octadecatrienoate) benzoyl bromide, 3,5-di-(9,12,15-octadecatrienoate) benzoyl iodide, α -linolenic: 3,5-di-(3,6,9-octadecatrienoate) benzoyl chloride, 3,5-di-(3,6,9-octadecatrienoate) benzoyl bromide, 3,5-di-(3,6,9-octadecatrienoate) benzoyl iodide, arachidonic: 3,5-di-(5,8,11,14-eicosatetraenoate) benzoyl chloride, 3,5-di-(5,8,11,14-eicosatetraenoate) benzoyl bromide, 3,5-di-(5,8,11,14-eicosatetraenoate) benzoyl iodide, eicosapentaenoic: 3,5-di-(5,8,11,14,17-eicosapentaenoate) benzoyl chloride, 3,5-di-(5,8,11,14,17-eicosapentaenoate) benzoyl bromide, 3,5-di-

(5,8,11,14,17-eicosapentaenoate) benzoyl iodide, docosahexanoic: 3,5-di-(4,7,10,13,16,19-docosahexaenoate) benzoyl chloride, 3,5-di-(4,7,10,13,16,19-docosahexaenoate) benzoyl bromide, and 3,5-di-(4,7,10,13,16,19-docosahexaenoate) benzoyl iodide.

5 In yet another aspect of the invention, a compound of Formula I may be made where each Y and each Z is independently selected from -O (ethers and heterocyclic rings), -O-C=O (esters), -O-C=O-O (carbonates), -O-C=O-NH, -O-C=O-NR, -NH-C=O-O, -NR-C=O-O (carbamates), -NH-C=O, -NR-C=O, -C=O-NH, -C=O-NR (primary and secondary amides) -NH, -NR (primary and secondary
10 amines), -N (heterocyclic rings), -S (heterocyclic rings), and halogen, each n and each m is independently the value of 0, 1, 2, 3, 4 or 5, where each A and each B is independently selected from H, R or absent, where each V and each W is independently selected from H, straight or branched alkyl of from 1 to 6 carbon atoms and cycloalkyl of from 3 to 8 carbon atoms, alkoxy, phenyl, benzyl or
15 halogen, and where R is independently selected from the group comprising alkyl with at least one carbon atom, aryl and aralkyl, and includes diastereoisomers of the foregoing.

Scheme 9 illustrates the general procedure for the preparation of the compounds of Formula I where each A and each B are independently selected from
20 the group comprising alkyl with at least one carbon atom, aryl, aralkyl, P_n, P_m (wherein P_n and P_m are each a protecting group), and H. It should be appreciated that with regard to compounds of Formulas IB, IC, ID, IE, and IF, A and B do not represent H; and, with regard to compounds of Formula IA and IE, when any Y and any Z is halogen, A or B is absent. In some embodiments A and B is alkyl with at
25 least two carbon atoms. In some embodiments the sum of m and n is greater than or equal to one but less than nine. In other embodiments the sum of m and n is three or more but less than nine.

Scheme 9.



Protection/Acylation of Benzoic Acid Halide Coupling Partner

- As depicted in Scheme 9, the benzoic acid derivative IA is optionally
- 5 acylated and/or optionally protected with a desired number and type of acyl group(s) equal to or less than n and/or with the desired number and type of protecting group(s) equal to or less than n to yield benzoic acid derivative IB where A is selected from the group comprising alkyl with at least one carbon atom, aryl, aralkyl, and P_n .
- 10 A variety of reaction pathways starting with benzoic acid derivative IA can be used to obtain the benzoic acid derivative IB. These include homogenous peracylation or polyacylation (condition I), homogenous polyprotection (condition II), selective acylation followed by selective protection (condition III), selective protection followed by selective acylation (condition IV), heterogeneous
- 15 peracylation (condition V), heterogeneous polyprotection (condition VI), selective acylation followed by homogeneous or heterogeneous polyprotection (condition VII) and selective protection followed by homogeneous or heterogeneous peracylation (condition IIX). It is contemplated that under condition VII, selective acylation may occur at one or more substituent sites and polyprotection may occur at
- 20 one or more substituents sites. In like manner, it is contemplated that under condition IIX, selective protection may occur at one or more substituent sites and peracylation may occur at one or more substituents sites. When each Y or each Z is halogen, the acylation and/or protection and diprotection steps may be entirely


omitted. Where one or more of Y and Z are halogen and one or more of Y and Z is neither H nor halogen, the acylation and/or protection and diprotection steps may be implemented.

It is to be understood from this description that benzoic acid IA includes
5 benzoic acid and monohydroxyl benzoic acids, dihydroxyl benzoic acids, trihydroxyl benzoic acids, tetrahydroxyl benzoic acids, and pentahydroxyl benzoic acids.

Benzoic acid IA also includes the monoamino benzoic acids 2-amino benzoic acid (anthranilic acid), 3-amino benzoic acid and 4-amino benzoic acid
10 (Vitamin H₁) in which case only conditions I or II can apply. Benzoic acid IA also includes but is not limited to the diamino benzoic acids 2,3-diamino benzoic acid, 2,4-diamino benzoic acid, 3,4-diamino benzoic acid, 3,5-diamino benzoic acid in which case conditions I–VI can apply. Benzoic acid IA also includes but is not limited to the monohydroxyl-monoamino benzoic acids. Illustrative examples of
15 which include but are not limited to 2-amino-5-hydroxy benzoic acid (5-hydroxyanthranilic acid), 3-amino-4-hydroxy benzoic acid, 4-amino-3-hydroxy benzoic acid, 3-amino-2-hydroxy benzoic acid (3-aminosalicylic acid), 4-amino-2-hydroxy benzoic acid (4-aminosalicylic acid) and 5-amino-2-hydroxybenzoic acid (5-aminosalicylic acid) in which case conditions I-VI can apply.

20 By way of illustration, the 3,5-diaminobenzoic acid analogs of IA, where $n=2$ and each Y is NH₂, the amines can be 1) both acylated with one or more acylating agents (condition I), 2) both protected with a one or more protecting agents (condition II), 3) selectively acylated with an acylating agent and selectively protected with a protecting agent (condition III), 4) selectively protected with a
25 protecting agent and selectively acylated with an acylating agent (condition IV), 5) heterogeneous diacylated with both amines acylated with two different acylating agents (condition V), or 6) heterogeneous diprotection with both amines protected with two different protecting agents (condition VI).


In the example of diacylation (condition I), both amines are reacted with at
30 least 2 equivalents of an acylating agent in a basic solvent (or in an inert solvent with added base) to give the diamide 3,5-diaminobenzoic acid derivative IB_a where A₁ and A₂ are the same. Preferably, a 5 molar equivalent excess of the acylating agent is used.



In the example of diprotection (condition II), both amines are reacted with at least 2 equivalents of a protection agent and under conditions known to one skilled in the art for the desired amine protecting group to give the diprotected 3,5-diaminobenzoic acid derivative IB_a where A₁ and A₂ are the same.

In the example of selective acylation followed by selective protection (condition III), 3,5-diaminobenzoic acid is first reacted with 1 equivalent or more of an acylating agent in a basic solvent (or in an inert solvent with added base) to give 3,5-diaminobenzoic acid derivative IB_b where NHA₁ represents an amide.

Intermediate IB_b is then reacted with 1 equivalent or more of a protection agent and under conditions known to one skilled in the art for the desired amine protecting group to give 3,5-diaminobenzoic acid derivative IB_a where NHA₂ represents a protected amine.



In the example of selective protection followed by selective acylation (condition IV), 3,5-diaminobenzoic acid is first reacted with 1 equivalent or more of a protection agent and under conditions known to one skilled in the art for the desired amine protecting group to give 3,5-diaminobenzoic acid derivative IB_b where NHA₁ represents a protected amine. Intermediate IB_b is then reacted with 1 equivalent or more of an acylating agent in a basic solvent (or in an inert solvent with added base) to give 3,5-diaminobenzoic acid derivative IB_a where NHA₂ represents an amide.

In the example of heterogeneous diacylation (condition V), 3,5-diaminobenzoic acid is first reacted with 1 equivalent or more of a first acylating agent in a basic solvent (or in an inert solvent with added base) to give 3,5-diaminobenzoic acid derivative IB_b where NHA₁ represents a first amide. Intermediate IB_b is then reacted with 1 equivalent or more of a second acylating

agent to give 3,5-diaminobenzoic acid derivative IB_a where NHA₂ represents a second amide.

In the example of heterogeneous diprotection (condition VI), 3,5-diaminobenzoic acid is first reacted with 1 equivalent or more of a first protection
5 agent under conditions known to one skilled in the art for the desired amine protecting group to give 3,5-diaminobenzoic acid derivative IB_b where NHA₁ represents a first protected amine. Intermediate IB_b is then reacted with 1 equivalent or more of a second protecting agent to give 3,5-diaminobenzoic acid derivative IB_a where NHA₂ represents a second protected amine.

10 Suitable acylating agents useful in the step or steps described above include those previously described for esterification. In the examples where acylation results in an amide (i.e. Y and/or Z is NH₂), acylation is followed by treatment with aqueous mild acid such as formic acid and optionally by recrystallization. Suitable protecting groups are described in T. W. Greene, Protective Groups in Organic
15 Synthesis, John Wiley & Sons, New York, N.Y., 1991, 2nd Edition.

Preferred amine protecting groups are those that can be selectively cleaved over a phenyl amine and include *tert*-butyloxycarbonyl (BOC), fluorinomethyloxycarbonyl (Fmoc), and benzyloxycarbonyl (Cbz).

Halogenation of Acid Chloride Coupling Partner

20 Returning to Scheme 9, benzoic acid derivative IB is reacted with a halogenating agent in an inert solvent to yield the benzoic acid halide derivative IC where X is any halogen.

Suitable halogenating agents useful in this step include but are not limited to any of the halogenating agents known to the art, such as thionyl halides including
25 thionyl chloride, thionyl bromide and thionyl iodide, oxalyl halides including oxalyl chloride, oxalyl bromide and oxalyl iodide, phosphorous tri- and pentahalides and phosphorous oxyhalides. Halogenation is carried out in an inert solvent including but not limited to halogenated alkanes such as methylene chloride and chloroform; cyclic ethers, such as THF and dioxane; N,N-dialkylsubstituted acylamides, such as
30 DMF and DMAC; and mixtures of said solvents.

The halogenation reaction may optionally include a buffering agent capable of associating with HX acid. Suitable buffering agents include weak bases.

Benzotriazole is a preferred buffering agent during halogenation because it precipitates as a salt after reacting with a HX acid.

The reaction is effected at a temperature from about 0 to about 100° C. Preferably, the reaction is effected using thionyl chloride or oxalyl chloride in a polar solvent such as methylene chloride, at 80°C, in the presence of benzotriazole to yield the corresponding acid chloride.

Optionally, the intermediate IC is recrystallized from hexane.

Preparation of Styrene Coupling Partner

Also depicted in Scheme 9 is styrene derivative ID. A variety of reaction pathways starting with styrene derivative ID can be used to obtain the styrene derivative IE. These pathways include homogenous peracylation or polyacylation (condition I), homogenous polyprotection (condition II), selective acylation followed by selective protection (condition III), selective protection followed by selective acylation (condition IV), heterogeneous peracylation (condition V), heterogeneous polyprotection (condition VI), selective acylation followed by homogeneous or heterogeneous polyprotection (condition VII) and selective protection followed by homogeneous or heterogeneous peracylation (condition IIX). It is contemplated that under condition VII, selective acylation may occur at one or more substituent sites and polyprotection may occur at one or more substituents sites. In like manner, it is contemplated that under condition IIX, selective protection may occur at one or more substituent sites and peracylation may occur at one or more substituents sites.

It is to be understood from this description that styrene derivative IA includes styrene and the monohydroxyl styrenes, dihydroxyl styrenes, trihydroxyl styrenes, tetrahydroxyl styrenes, and pentahydroxyl styrenes.

Styrene derivative ID also includes the monoamino styrenes 2-amino styrene, 3-amino styrene and 4-amino styrene in which case only conditions I or II can apply. Styrene derivative ID also includes but is not limited to the diamino styrenes 2,3-diamino styrene, 2,4-diamino styrene, 3,4-diamino styrene, 3,5-diamino styrene in which case conditions I–VI can apply.

Styrene derivative ID also includes but is not limited to the monohydroxyl-monoamino styrenes. Illustrative examples of which include but are not limited to 2-amino-5-hydroxy styrene, 3-amino-4-hydroxy styrene, 4-amino-3-hydroxy

styrene, 3-amino-2-hydroxy styrene, 4-amino-2-hydroxy styrene and 5-amino-2-hydroxystyrene in which case conditions I-VI can apply.

By way of illustration, the monoamino styrene analogs of ID (where $n=1$ and Z is NH_2) has one amine that can be 1) acylated with an acylating agent (condition I) or 2) protected with a protecting agent (condition II).

Coupling Step via Decarbonylative Heck Reaction

Returning to Scheme 9, benzoic acid halide IC is reacted with styrene IE via a decarbonylative coupling reaction to give stilbene IF. The reaction conditions associated with the coupling reaction can include an inert apolar, high-boiling point aromatic or hydrocarbon solvent with a transition metal catalyst, N-heterocyclic carbene-type ligand and base.

Suitable inert solvents useful in the coupling step include but are not limited to benzene, toluene, xylene, chlorobenzene, decane, dodecane, and mixtures of those solvents. Xylene is a preferred solvent in the coupling step.

A preferred transition metal catalyst used in the coupling step is $\text{Pd}(\text{OAc})_2$.

A preferred N-heterocyclic carbene-type ligands useful in the coupling step is N,N-bis-(2,6-diisopropylphenyl) 4,5-dihydroimidazolium chloride in an amount of 0.1 – 10 mol %, preferably 5-10 mol %.

Non-coordinating amine bases result in greater yields than coordinating amine bases. Examples of non-coordinating bases useful in the process include but are not limited to N-ethylmorpholine (NEM), N,N-dimethylbenzylamine, dimethylaminopyridine (DMAP), proton sponge (1,8-diaminonaphthalene), diisopropylethylamine (Hunig's base) and triethylamine. N-ethylmorpholine and N,N-dimethylbenzylamine are preferred bases in the coupling step. Use of smaller amines capable of transition metal coordination still work to obtain the coupled product IF but are less efficient. Furthermore, phosphine ligands can also be added to the reaction; however, added phosphine ligand inhibits the reaction giving greatly lowered yields.

The reaction is effected at a temperature from about 0 to about 150° C. Preferably, the reaction is refluxed in xylene.

The decarbonylative Heck reaction depicted in Scheme 9 is concluded using standard work-up conditions and the reaction products can be purified using a

variety of techniques known to those skilled in the art including silica gel chromatography.

Deprotection Step


After performing the coupling reaction, in those cases where one or more protecting groups exist on the stilbene derivative IF, the protection groups may be deprotected as described in T. W. Greene, Protective Groups in Organic Synthesis, John Wiley & Sons, New York, N.Y., 1991, 2nd Edition for each of the respective protection groups and may require one or more steps to complete.

In the example of stilbene derivatives IF akin to resveratrol, where $Y=NH_2$ at the 3 and 5 positions, $n=2$, $Z=NH_2$ at the 4' position and $m=1$, the derivative may be 1) deacylated at all three amine sites (3,5, and 4'), 2) deprotected and/or deacylated at all three amine sites and 3) selectively deprotected at any of the three amine sites.

Olefin Isomerization

As discussed previously, the stilbene compounds of the present invention possess a conjugated double bond. The double bond may exist in the *cis* or *trans* configuration. When V and W are H, the decarbonylative Heck reaction yields a *trans* product. However, depending on the nature of the V and W substitutions (i.e. other than H), the decarbonylative Heck reaction of the present invention may yield a *cis* or *trans* product. The present invention includes both *cis* and *trans* isomers as mixtures or in purified form. Furthermore, one aspect of the present invention includes a process for isomerizing between the *cis* and *trans* configurations. As depicted in Scheme 10, the isomerization step may be conducted to the stilbene derivatives when Y is acylated, protected, or hydrolyzed. Compounds of Formula I can be diluted in an inert solvent and irradiated with ultraviolet light, preferably of wavelength $\lambda = 280-350$ nm. This ultraviolet irradiation is carried in organic solution, at temperatures of 0° to 50° , preferably of 10° to 30° . The irradiation can be carried out with the aid of the most diverse commercial ultraviolet apparatuses. By way of example, irradiation can be carried out with a medium-pressure mercury vapor lamp through a pyrex filter. The irradiation time is generally from 2 to 100 hours, preferably 5 to 50 hours, especially 10 to 20 hours.

Scheme 10.



Compounds of Formulas I and 1-7 are capable of further forming both pharmaceutical and cosmetically acceptable formulations.

Pharmaceutical Uses

5 The compounds of the present invention can be prepared and administered in a wide variety of oral and parenteral dosage forms. Thus, the compounds of the present invention can be administered by injection, that is, intravenously, intramuscularly, intracutaneously, subcutaneously, intraduodenally, or intraperitoneally. Also, the compounds of the present invention can be administered
10 by inhalation, for example, intranasally. Additionally, the compounds of the present invention can be administered transdermally. It will be obvious to those skilled in the art that the following dosage forms may comprise as the active component, either a compound of Formula 1 or a corresponding pharmaceutically acceptable salt of a compound of Formula 1.

15 For preparing pharmaceutical compositions from the compounds of the present invention, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier can be one or more substances which may also act as diluents, flavoring agents, binders, preservatives,
20 tablet disintegrating agents, or an encapsulating material.

In powders, the carrier is a finely divided solid which is in a mixture with the finely divided active component.

In tablets, the active component is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape and
25 size desired.

The powders and tablets preferably contain from five or ten to about seventy percent of the active compound. Suitable carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter,

and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as a carrier providing a capsule in which the active component with or without other carriers, is surrounded by a carrier, which is thus in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid dosage forms suitable for oral administration.

For preparing suppositories, a low melting wax, such as a mixture of fatty acid glycerides or cocoa butter, is first melted and the active component is dispersed homogeneously therein, as by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool, and thereby to solidify.

Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water propylene glycol solutions. For parenteral injection liquid preparations can be formulated in solution in aqueous polyethylene glycol solution.

Aqueous solutions suitable for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizing and thickening agents as desired.

Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-known suspending agents.

Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. These preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

The pharmaceutical preparation is preferably in unit dosage form. In such form the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, table, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

The quantity of active component in a unit dose preparation may be varied or adjusted from 0.1 mg to 1000 mg preferably 0.5 mg to 100 mg according to the particular application and the potency of the active component. The composition can, if desired, also contain other compatible therapeutic agents.

5 Cosmetic Uses

For preparing cosmetic compositions from the processes and compounds of the present invention, cosmetically acceptable vehicles or carriers facilitate distribution when the composition is applied to the skin. The vehicle may be aqueous, anhydrous or an emulsion. Preferably, the compositions are aqueous or an emulsion, especially water-in-oil or oil-in-water emulsion. Water when present will
10 be in amounts which may range from 5 to 99%, preferably from 20 to 70%, optimally between 35 and 60% by weight.

Besides water, relatively volatile solvents may also serve as carriers within compositions of the present invention. Most preferred are monohydric C₁-C₃
15 alkanols. These include ethyl alcohol, methyl alcohol and isopropyl alcohol. The amount of monohydric alkanol may range from 1 to 70%, preferably from 10 to 50%, optimally between 15 to 40% by weight.

Emollient materials may also serve as cosmetically acceptable carriers. These may be in the form of silicone oils and synthetic esters. Amounts of the
20 emollients may range anywhere from 0.1 to 50%, preferably between 1 and 20% by weight.

Silicone oils may be divided into the volatile and non-volatile variety. The term "volatile" as used herein refers to those materials which have a measurable vapor pressure at ambient temperature. Volatile silicone oils are preferably chosen
25 from cyclic or linear polydimethylsiloxanes containing from 3 to 9, preferably from 4 to 5, silicon atoms. Linear volatile silicone materials generally have viscosities less than about 5 centistokes at 25° C. while cyclic materials typically have viscosities of less than about 10 centistokes. Nonvolatile silicone oils useful as an emollient material include polyalkyl siloxanes, polyalkylaryl siloxanes and
30 polyether siloxane copolymers. The essentially non-volatile polyalkyl siloxanes useful herein include, for example, polydimethyl siloxanes with viscosities of from about 5 to about 25 million centistokes at 25° C. Among the preferred non-volatile

emollients useful in the present compositions are the polydimethyl siloxanes having viscosities from about 10 to about 400 centistokes at 25° C.

Among the ester emollients are: (1) Alkenyl or alkyl esters of fatty acids having 10 to 20 carbon atoms. Examples thereof include isoarachidyl neopentanoate, isononyl isononanoate, oleyl myristate, oleyl stearate, and oleyl oleate; (2) ether-esters such as fatty acid esters of ethoxylated fatty alcohols; (3) polyhydric alcohol esters. Ethylene glycol mono and di-fatty acid esters, diethylene glycol mono- and di-fatty acid esters, polyethylene glycol (200-6000) mono- and di-fatty acid esters, propylene glycol mono- and di-fatty acid esters, polypropylene glycol 2000 monooleate, polypropylene glycol 2000 monostearate, ethoxylated propylene glycol monostearate, glyceryl mono- and di-fatty acid esters, polyglycerol poly-fatty esters, ethoxylated glyceryl mono-stearate, 1,3-butylene glycol monostearate, 1,3-butylene glycol distearate, polyoxyethylene polyol fatty acid ester, sorbitan fatty acid esters, and polyoxyethylene sorbitan fatty acid esters are satisfactory polyhydric alcohol esters; (4) wax esters such as beeswax, spermaceti, myristyl myristate, stearyl stearate and arachidyl behenate; and, (5) sterols esters, of which cholesterol fatty acid esters are examples thereof; fatty acids having from 10 to 30 carbon atoms may also be included as cosmetically acceptable carriers for compositions of this invention. Illustrative of this category are pelargonic, lauric, myristic, palmitic, stearic, isostearic, hydroxystearic, oleic, linoleic, ricinoleic, arachidic, behenic and erucic acids.

Humectants of the polyhydric alcohol-type may also be employed as cosmetically acceptable carriers in compositions of this invention. The humectant aids in increasing the effectiveness of the emollient, reduces scaling, stimulates removal of built-up scale and improves skin feel. Typical polyhydric alcohols include glycerol, polyalkylene glycols and more preferably alkylene polyols and their derivatives, including propylene glycol, dipropylene glycol, polypropylene glycol, polyethylene glycol and derivatives thereof, sorbitol, hydroxypropyl sorbitol, hexylene glycol, 1,3-butylene glycol, 1,2,6-hexanetriol, ethoxylated glycerol, propoxylated glycerol and mixtures thereof. For best results the humectant is preferably propylene glycol or sodium hyaluronate. The amount of humectant may range anywhere from 0.5 to 30%, preferably between 1 and 15% by weight of the composition.

Thickeners may also be utilized as part of the cosmetically acceptable carrier of compositions according to the present invention. Typical thickeners include crosslinked acrylates (e.g. Carbopol 982), hydrophobically-modified acrylates (e.g. Carbopol 1382), cellulosic derivatives and natural gums. Among useful cellulosic derivatives are sodium carboxymethylcellulose, hydroxypropyl methylcellulose, hydroxypropyl cellulose, hydroxyethyl cellulose, ethyl cellulose and hydroxymethyl cellulose. Natural gums suitable for the present invention include guar, xanthan, sclerotium, carrageenan, pectin and combinations of these gums. Amounts of the thickener may range from 0.0001 to 5%, usually from 0.001 to 1%, optimally from 0.01 to 0.5% by weight.

Collectively the water, solvents, silicones, esters, fatty acids, humectants and/or thickeners will constitute the cosmetically acceptable carrier in amounts from 1 to 99.9%, preferably from 80 to 99% by weight.

An oil or oily material may be present, together with an emulsifier to provide either a water-in-oil emulsion or an oil-in-water emulsion, depending largely on the average hydrophilic-lipophilic balance (HLB) of the emulsifier employed.

Surfactants may also be present in cosmetic compositions of the present invention. Total concentration of the surfactant will range from 0.1 to 40%, preferably from 1 to 20%, optimally from 1 to 5% by weight of the composition. The surfactant may be selected from the group consisting of anionic, nonionic, cationic and amphoteric actives. Particularly preferred nonionic surfactants are those with a C₁₀-C₂₀ fatty alcohol or acid hydrophobe condensed with from 2 to 100 moles of ethylene oxide or propylene oxide per mole of hydrophobe; C₂-C₁₀ alkyl phenols condensed with from 2 to 20 moles of alkylene oxide; mono- and di-fatty acid esters of ethylene glycol; fatty acid monoglyceride; sorbitan, mono- and di-C₈-C₂₀ fatty acids; block copolymers (ethylene oxide/propylene oxide); and polyoxyethylene sorbitan as well as combinations thereof. Alkyl polyglycosides and saccharide fatty amides (e.g. methyl gluconamides) are also suitable nonionic surfactants.

Preferred anionic surfactants include soap, alkyl ether sulfate and sulfonates, alkyl sulfates and sulfonates, alkylbenzene sulfonates, alkyl and dialkyl sulfosuccinates, C₈-C₂₀ acyl isethionates, acyl glutamates, C₈-C₂₀ alkyl ether phosphates and combinations thereof.

Sunscreens may be present in cosmetic compositions of the present invention. Sunscreens include those materials commonly employed to block ultraviolet light. Illustrative compounds are the derivatives of PABA, cinnamate and salicylate. For example, avobenzophenone (Parsol 1789.RTM.) octyl methoxycinnamate and 2-hydroxy-4-methoxy benzophenone (also known as oxybenzone) can be used. Octyl methoxycinnamate and 2-hydroxy-4-methoxy benzophenone are commercially available under the trademarks, Parsol MCX and Benzophenone-3, respectively. The exact amount of sunscreen employed in the compositions can vary depending upon the degree of protection desired from the sun's UV radiation.

Many cosmetic compositions, especially those containing water, must be protected against the growth of potentially harmful microorganisms. Preservatives are, therefore, often necessary. Suitable preservatives include alkyl esters of p-hydroxybenzoic acid, hydantoin derivatives, propionate salts, and a variety of quaternary ammonium compounds. Particularly preferred preservatives of this invention are methyl paraben, propyl paraben, phenoxyethanol and benzyl alcohol. Preservatives will usually be employed in amounts ranging from about 0.1% to 2% by weight of the composition.

Powders may be incorporated into the cosmetic composition of the invention. These powders include chalk, talc, Fullers earth, kaolin, starch, smectites clays, chemically modified magnesium aluminum silicate, organically modified montmorillonite clay, hydrated aluminum silicate, fumed silica, aluminum starch octenyl succinate and mixtures thereof.

The composition can be packaged in a suitable container to suit its viscosity and intended use by the consumer. For example, a lotion or fluid cream can be packaged in a bottle or a roll-ball applicator, or a capsule, or a propellant-driven aerosol device or a container fitted with a pump suitable for finger operation. When the composition is a cream, it can simply be stored in a non-deformable bottle or squeeze container, such as a tube or a lidded jar.

Cosmetic formulations according to the invention are used in various fields. For example, the following preparations especially come into consideration: skin-care preparations, e.g. skin-washing and cleansing preparations in the form of tablet-form or liquid soaps, synthetic detergents or washing pastes; bath preparations, e.g.

liquid (foam baths, milks, shower preparations) or solid bath preparations, e.g. bath cubes and bath salts; skin-care preparations, e.g. skin emulsions, multi-emulsions or skin oils; cosmetic personal care preparations, e.g. facial make-up in the form of day creams or powder creams, face powder (loose or pressed), rouge or cream make-up, eye-care preparations, e.g. eyeshadow preparations, mascara, eyeliner, eye creams or eye-fix creams; lip-care preparations, e.g. lipstick, lip gloss, lip contour pencils, nail-care preparations, such as nail varnish, nail varnish remover, nail hardeners, or cuticle removers; intimate hygiene preparations, e.g. intimate washing lotions or intimate sprays; foot-care preparations, e.g. foot baths, foot powders, foot creams or foot balsams, special deodorants and antiperspirants or callous-removing preparations; light-protective preparations, such as sun milks, lotions, creams, oils, sun blocks or tropicals, pre-tanning preparations or after-sun preparations; skin-tanning preparations, e.g. self-tanning creams; depigmenting preparations, e.g. preparations for bleaching the skin or skin-lightening preparations; insect-repellents, e.g. insect-repellent oils, lotions, sprays or sticks; deodorants, such as deodorant sprays, pump-action sprays, deodorant gels, sticks or roll-ons; antiperspirants, e.g. antiperspirant sticks, creams or roll-ons; preparations for cleansing and caring for blemished skin, e.g. synthetic detergents (solid or liquid), peeling or scrub preparations or peeling masks; hair-removal preparations in chemical form (depilation), e.g. hair-removing powders, liquid hair-removing preparations, cream- or paste-form hair-removing preparations, hair-removing preparations in gel form or aerosol foams; shaving preparations, e.g. shaving soap, foam shaving creams, non-foaming shaving creams, foams, gels, preshave preparations for dry shaving, aftershaves or aftershave lotions; fragrance preparations, e.g. fragrances (eau de Cologne, eau de toilette, eau de parfum, parfum de toilette, perfume), perfume oils or perfume creams; dental-care, denture-care and mouth-care preparations, e.g. toothpastes, gel tooth-pastes, tooth powders, mouthwash concentrates, anti-plaque mouthwashes, denture cleaners or denture fixatives; and, cosmetic hair-treatment preparations, e.g. hair-washing preparations in the form of shampoos and conditioners, hair-care preparations, e.g. pre-treatment preparations, hair tonics, styling creams, styling gels, pomades, hair rinses, treatment packs, intensive hair treatments, hair-structuring preparations, e.g. hair-waving preparations for permanent waves (hot wave, mild wave, cold wave), hair-straightening preparations,

liquid hair-setting preparations, hair foams, hair sprays, bleaching preparations, e.g. hydrogen peroxide solutions, lightening shampoos, bleaching creams, bleaching powders, bleaching pastes or oils, temporary, semi-permanent or permanent hair colorants, preparations containing self-oxidizing dyes, or natural hair colorants, such as henna or camomile.

The compounds of the present invention have significantly improved biological activity and can be stored over a long period without alteration.

The following non-limiting examples illustrate the preferred methods for preparing the compounds of the invention.

10 Nutritional Uses


When the composition is incorporated into various media such as foods, it may simply be orally ingested. The food can be a dietary supplement (such as a snack or wellness dietary supplement) or, especially for animals, comprise the nutritional bulk (e.g., when incorporated into the primary animal feed).

15 Examples

Air and moisture sensitive reagents were introduced via dry syringe or cannula. Toluene, xylene, pyridine, ethyl acetate, and N-methyl morpholine were distilled from CaH₂. DMF were dried by storage over 4Å Molecular sieves. Reagents were purchased from Aldrich and Lancaster. Flash chromatography was carried out using 60-230 mesh silica gel. Radical chromatography was performed using 1, 2, and 4 mm plates loaded with 230-400 mesh PF-254 gypsum bound silica. Analytical thin-layer chromatography (TLC) was performed with Merck silica gel 60 F₂₅₄, 0.25 mm pre-coated TLC plates. TLC plates were visualized using UV₂₅₄. All ¹H NMR spectra were obtained with 300 Varian spectrometers using TMS (0.00ppm), Chloroform (7.26 ppm), or acetone-d₆ (2.05 ppm) as an internal reference. Signals are reported as m (multiplet), s (singlet), d (doublet), t (triplet), q (quartet), and bs (broad singlet). ¹³C NMR were obtained with 75 MHz Varian spectrometer using TMS (0.0 ppm), Chloroform (77.2 ppm), or acetone-d₆ (30.6 ppm) as the internal standard. Mass spectra data (HRMS, EI) were obtained from the Brigham Young University mass spectrometry facility. Combustion analysis was performed by M-H-W Laboratories, Phoenix, AZ.

The following are prepared according to Scheme 11.

Scheme 11.



5 1. Preparation of 3,5-bis(methoxymethoxy)benzoic acid.

A flame dried flask was charged with dry DMF (75mL) and 60% oil dispersion NaH (3.8 g, 95 mmol). A solution of 3,5-dihydroxybenzoic acid (4.6 g, 30 mmol) in DMF (25mL) was added dropwisely over 20 minutes. The mixture was allowed to stir for one hour under N₂. MOMCl (7.5 mL, 100 mmol) was added slowly so that the inner temperature did not exceed 50°C. After 30 hours, the insoluble material was filtered off and the filtrate was concentrated to an oil residue, which was partitioned between benzene and water. The water layer was extracted with benzene for another three times. The combined benzene extracts was dried over Na₂SO₄ and concentrated to pale yellow oil, which was dissolved in 50 mL methanol. 2N Aqueous NaOH (25 mL, 50 mmol) was added and the mixture was stirred for three hours. The mixture was concentrated and dissolved in 30mL water. The aqueous solution was washed with benzene and acidified with 10% aqueous HCl. The precipitated white solid was filtered and washed with water and dried to give 6.6g (91%) product, which can be further purified by recrystallization from EtOAc-hexane. Data are: ¹H NMR (CDCl₃, 300 MHz) δ 7.44 (d, 2H), 6.98 (t, 1H), 5.21 (s, 4H), 3.50 (s, 6H); ¹³C NMR (CDCl₃, 75 MHz) δ 170.9, 158.4, 131.4, 111.5,

110.7, 94.7, 56.4; mp = 129-130 °C; HRMS (EI⁺) found 242.0796 M⁺, calcd 242.0790 for C₁₁H₁₄O₆.

2. Preparation of 3,5-bis(methoxymethoxy)benzoyl chloride.

A stock solution was prepared by dissolving benzotriazole (1.49 g, 0.0125 mmol), thionyl chloride (0.91 mL, 0.0125 mmol) in 8.0 mL DCM. Reaction was carried out by adding the stock solution intermittently into a stirred solution of 3,5-dimethoxybenzoic acid (2.42 g, 10 mmol) in 200 mL DCM. Before the addition was complete, benzotriazole hydrochloride started precipitating out as white solid. The mixture was stirred for another ten minutes. After filtration, the filtrate was stirred with MgSO₄·7H₂O (5 g) to destroy excess thionyl chloride. The white solid was filtered off and the filtrate was concentrated to give 2.5 g (97%) crude product, which was used for the next step without further purification. Data are: ¹H NMR (CDCl₃, 300 MHz) δ 7.44 (d, 2H), 7.04 (t, 1H), 5.20 (s, 4H), 3.49 (s, 6H); ¹³C NMR (CDCl₃, 75 MHz) δ 168.1, 158.5, 135.3, 112.6, 111.9, 94.7, 56.4; HRMS (EI⁺) found 260.0465 M⁺, calcd 260.0452 for C₁₁H₁₃O₅Cl.

3. Preparation of 4'-acetoxy-3,5-bis(methoxymethoxy)stilbene.

A 50 mL round bottom flask was charged with p-xylene (20 mL), Pd II catalyst (22.5 mg, 0.1 mmol), 1,3-bis-(2,6-diisopropylphenyl) imidazolinium chloride (42.7 mg, 0.1 mmol), 3,5-bis(methoxymethoxy)benzoyl chloride (2.42 g, 10 mmol), 4-acetoxystyrene (1.94 g, 12 mmol), and N-methyl morpholine (1.38 g, 12 mmol). The mixture was stirred at 120 °C for 3.5 h under nitrogen atmosphere. Then it was cooled to room temperature and EtOAc was added and filtered. The filtrate was washed with brine and dried over Na₂SO₄. Then it was filtered and purified via flash chromatography and gave the product (2.1 g, 59%) as white solid. Data are: ¹H NMR (CDCl₃, 300 MHz) δ 7.48 (d, 2H), 7.08-6.93 (m, 4H), 6.86 (d, 2H), 6.66 (t, 1H), 5.19 (s, 4H), 3.50 (s, 6H), 2.30 (s, 3H); ¹³C NMR (CDCl₃, 75 MHz) δ 169.5, 158.7, 150.3, 139.5, 135.0, 128.7, 128.5, 127.6, 121.9, 108.0, 104.5, 94.6, 56.2, 21.2; HRMS (EI⁺) found 358.1409 M⁺, calcd 358.1416 for C₂₀H₂₂O₆.

4. Preparation of 4'-acetoxy-3,5-dihydroxystilbene.


To a solution of 4'-acetoxy-3,5-bis(methoxymethoxy)stilbene (0.358 g, 1 mmol) in dry DCM (50 mL) and dry CH₃CN (50 mL) were added NaI (1.8 g, 24 mmol) and freshly distilled TMSCl (1.52 g, 24 mmol). The mixture was stirred under

argon for 15 minutes. The solution was diluted with DCM (50mL) and washed with a fresh aqueous saturated solution of $\text{Na}_2\text{S}_2\text{O}_3$ ($3 \times 40\text{mL}$) and saturated NaHCO_3 , and water. The organic layer was dried over Na_2SO_4 , filtered and concentrated. The crude product was purified by flash column and gave 0.20 g product (72%). Data are: ^1H NMR (Aceton- d_6 , 300 MHz) δ 8.25 (s, 1H), 7.60 (m, 2H), 7.13-7.08 (m, 4H), 6.59 (d, 2H), 6.32 (t, 1H), 2.25 (s, 3H); ^{13}C NMR (Aceton- d_6 , 75 MHz) δ 169.7, 159.7, 151.4, 140.4, 136.0, 130.0, 128.3, 128.3, 123.0, 106.1, 103.3, 21.1; HRMS (EI^+) found 270.0889 M^+ , calcd 270.0892 for $\text{C}_{16}\text{H}_{16}\text{O}_4$.

The following are prepared according to Scheme 12.

10

Scheme 12.



5. Preparation of 3,5-diacetoxybenzoic acid.

15

To a solution of 3,5-dihydroxybenzoic acid (7.71 g, 50 mmol) in EtOAc (110 mL) were added acetic anhydride (12.25 mL, 130 mmol) and pyridine (8.08 mL, 100mmol) in an ice-water bath under a nitrogen atmosphere. The mixture was stirred at 0°C for 40 min and then stirred at ambient temperature for 4 h. 98% Formic acid (2.36 mL, 60 mmol) was added and the resulting mixture was stirred for 1 h. Then, it

was poured into water and extracted with EtOAc. The organic phase was washed with water and brine, dried over Na₂SO₄, filtered, and evaporated in vacuo. Purification of the residue via recrystallization from n-heptane/EtOAc provided 3,5-diacetoxybenzoic acid (8.97 g, 75.3%) as white solid. Data are: ¹H NMR (CDCl₃, 300 MHz) δ 12.1 (bs, 1H), 7.73 (d, 2H), 7.21 (t, 1H), 2.32 (s, 6H); ¹³C NMR (CDCl₃, 75 MHz) δ 170.4, 169.0, 151.2, 131.5, 121.3, 121.0, 21.2; mp = 161-162 °C; HRMS (EI⁺) found 238.0475 M⁺, calcd 238.0477 for C₁₁H₁₀O₆; Anal. Calcd for C₁₁H₁₀O₆: C, 55.47; H, 4.23. Found: C, 55.62; H, 4.37.

6. Preparation of 3,5-diacetoxybenzoyl chloride.

To a mixture of 3,5-diacetoxybenzoic acid (8.00 g, 33.59 mmol) and dimethylformamide (5 drops) in toluene was added freshly distilled thionylchloride (16 mL). The mixture was stirred for 15 min under nitrogen atmosphere at ambient temperature. Then it was refluxed for 2 h at 80 °C in a hot water bath. The excess thionyl chloride was evaporated in vacuo and toluene was added. Insoluble yellow solid was discarded. Toluene was evaporated in vacuo and gave 3,5-diacetoxybenzoyl chloride (8.23g, 95.5%), which can be recrystallized from hexane. Data are: ¹H NMR (CDCl₃, 300 MHz) δ 7.75 (d, 2H), 7.29 (t, 1H), 2.34 (s, 6H); ¹³C NMR (CDCl₃, 75 MHz) δ 168.8, 167.0, 151.4, 135.3, 122.8, 122.0, 21.2; mp = 89.5-91 °C; HRMS (EI⁺) found 256.0130 M⁺, calcd 256.0139 for C₁₁H₉O₅Cl; Anal. Calcd for C₁₁H₉O₅Cl: C, 51.48; H, 3.53. Found: C, 51.60; H, 3.68.

7. Preparation of 4-chloroacetoxystyrene.

A flask was charged with 4-acetoxystyrene (10 g, 61.73 mmol), methanol (30mL), and KOH (0.125 g, 2.23mmol), and 1 drop of water. After the mixture was stirred for 5 minutes under N₂, the temperature was raised to 65 °C. The mixture was stirred for 1.5 h and then cooled to room temperature. Acetic acid (0.144 g, 2.44mmol) in methanol (0.5mL) was added slowly over 5 min. The mixture was stirred for another 5 min and then concentrated. The residue was dissolved in toluene and filtered. The filtrate was cooled to -78 °C and the 4-hydroxystyrene was precipitated, filtered and dried to give 4.5 g product (60%).

To a solution of 4-hydroxystyrene (1.22 g, 10 mmol) in ethyl ether 140 mL was added chloroacetic chloride (2.26 g, 20 mmol) and Et₃N (1.62 g, 16 mmol). After it was stirred for 2 h, the mixture was washed with brine and water. The ether

solution was dried over Na_2SO_4 and concentrated. After purification by flash column, the 4-chloroacetoxystyrene (1.8 g, 92%) was obtained. Data are: ^1H NMR (CDCl_3 , 300 MHz) δ 7.46 (d, 2H, $J=1.8$ Hz), 7.11 (d, 2H, $J=8.4$ Hz), 6.73 (dd, 1H, $J=17.7$ Hz), 5.74 (d, 1H, $J=18$ Hz), 5.29 (d, $J=11.1$ Hz), 4.31 (s, 2H). ^{13}C NMR (CDCl_3 , 75 MHz) δ 166.0, 150.0, 136.0, 135.8, 127.4, 121.3, 114.6, 41.0.

8. Preparation of 3,5-diacetoxy-4'-chloroacetoxy stilbene.


A 10 mL round bottom flask was charged with p-xylene (4 mL), Pd II catalyst (4.5 mg, 0.02 mmol), 1,3-bis-(2,6-diisopropylphenyl) imidazolinium chloride (8.6 mg, 0.02 mmol), 3,5-diacetoxybenzoyl chloride (0.7695 g, 3 mmol), 4-chloroacetoxystyrene (0.393 g, 2 mmol), and N-methyl morpholine (0.276 g, 2.4 mmol). The mixture was stirred at 120 °C for 3.5 h under nitrogen atmosphere. Then it was cooled to room temperature and EtOAc was added and filtered. The filtrate was washed with brine and dried over Na_2SO_4 . Then it was filtered and purified via flash chromatography and gave the product (0.54 g, 69.4%) as white solid. Data are: ^1H NMR (CDCl_3 , 300 MHz) δ 7.47 (d, 2H, $J=8.7$ Hz), 7.14-7.10 (m, 4H), 6.99 (q, 2H, $J=16.2$ Hz), 6.83 (t, 1H, $J=2.1$ Hz), 4.29 (s, 2H), 2.29 (s, 6H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 169.1, 165.9, 151.4, 150.1, 139.5, 135.1, 129.5, 127.9, 127.7, 121.6, 117.1, 114.7, 41.0, 21.2; HRMS (EI^+) found 388.0710 M^+ , calcd 388.0714 for $\text{C}_{20}\text{H}_{17}\text{ClO}_6$.

9. Preparation of 3,5-diacetoxy-4'-hydroxy stilbene.

A solution of 3,5-diacetoxy-4'-chloroacetoxy stilbene (0.388 g, 1 mmol) in 50% aqueous pyridine, which was adjusted to pH 6.7 with hydrochloric acid, was stirred for 6 h at room temperature. The mixture was concentrated and diluted with EtOAc. The mixture was washed with 1 N aqueous HCl, saturated NaHCO_3 and water. Then the EtOAc solution was dried and concentrated. After purification by radial chromatography, the product (0.28 g, 90%) was obtained. Data are: ^1H NMR (Aceton- d_6 , 300 MHz) δ 8.55 (bs, 1H), 7.47 (m, 2H), 7.25-7.00 (m, 4H), 6.86 (m, 2H), 6.82 (t, 1H, $J=2.1$ Hz), 2.27 (s, 3H); ^{13}C NMR (Aceton- d_6 , 75 MHz) δ 169.5, 158.7, 152.7, 141.2, 131.4, 129.5, 129.2, 124.8, 117.5, 116.6, 115.1, 21.0; HRMS (EI^+) found 312.0993 M^+ , calcd 312.0998 for $\text{C}_{18}\text{H}_{16}\text{O}_5$.

The following are prepared according to Scheme 13.

Scheme 13.



5 10. Preparation of 3-acetoxy-5-hydroxybenzoic acid.

To 3,5-dihydroxybenzoic acid (20 g, 129 mmol) was added NaOH (15.6 g, 390 mmol) in 100 mL water. After the mixture was cooled to 0 °C, Ac₂O (13.2 g, 129 mmol) was added. The solution was stirred for 40 min and then acidified with 10% H₂SO₄ at 0 °C. The precipitate was filtered and washed with cold water. The crude product was recrystallized from water to give 3-acetoxy-5-hydroxybenzoic acid (15.2 g, 60%). Data are: ¹H NMR (Aceton-d₆, 300 MHz) δ 8.96 (s, 1H), 7.40 (dd, 1H, J=1.5Hz), 7.26 (dd, 1H, J=1.5Hz), 6.86 (t, 1H, 2.1H), 2.27 (s, 3H); HRMS (EI⁺) found 196.0380 M⁺, calcd 196.0372 for C₉H₈O₅.

15 11. Preparation of 3-acetoxy-5-levulinoxybenzoic acid.

To a solution of 3-acetoxy-5-hydroxybenzoic acid (2.45 g, 12.5 mmol) in CH₂Cl₂ (25 mL) was added levulinic anhydride (5.4 g, 25 mmol) and pyridine (1.22 mL, 15 mmol) at 0 °C. The mixture was stirred at 0 °C for another 0.5 h and then warmed to room temperature. After 3 h, 98% formic acid was added and stirred for 1

h. The mixture was washed with 1 N HCl, saturated NaHCO₃ and water. Then it was dried by Na₂SO₄ and concentrated. The crude product was purified by flash column and give 3-acetoxy-5-levulinoxybenzoic acid (3.52 g, 96%). Data are: ¹H NMR (CDCl₃, 300 MHz) δ 7.72 (d, 2H, J=1.2Hz), 7.20 (t, 1H, 1.2Hz), 2.85 (m, 4H), 2.32 (s, 3H), 2.24 (s, 3H); ¹³C NMR (CDCl₃, 75 MHz) δ 206.6, 177.6, 171.0, 170.0, 169.0, 151.2, 131.7, 121.1, 121.1, 121.0, 38.1, 30.0, 28.3, 21.2; HRMS (EI⁺) found 294.0741 M⁺, calcd 294.0740 for C₁₄H₁₄O₇.

12. Preparation of 3-acetoxy-5-levulinoxybenzoyl chloride.

A stock solution was prepared by dissolving benzotriazole(1.49 g, 12.5 mmol), thionyl chloride(0.91mL, 12.5mmol) in 8.0mL DCM. Reaction was carried out by adding the stock solution intermittently into a stirred solution of 3-acetoxy-5-levulinoxybenzoic acid (2.94 g, 10 mmol) in 200ml DCM. Before the addition was complete, benzotriazole hydrochloride started precipitating out as white solid. The mixture was stirred for another ten minutes. After filtration, the filtrate was stirred with MgSO₄·7H₂O (5g) to destroy the excess thionyl chloride. After concentration, the residue was extracted by hot dry hexane several times and recrystallized from hexane to give 3-acetoxy-5-levulinoxybenzoyl chloride (1.4 g, 45%). Data are: ¹H NMR (CDCl₃, 300 MHz) δ 7.75 (d, 2H, J=2.1Hz), 7.28 (t, 1H, J=2.1Hz), 2.86 (m, 4H), 2.33 (s, 3H), 1.24 (s, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 206.3, 170.9, 168.7, 167.1, 151.42, 151.38, 135.3, 122.8, 122.1, 121.9, 38.0, 30.0, 28.3, 21.2; HRMS (EI⁺) found 312.0390 M⁺, calcd 312.0401 for C₁₄H₁₃O₆Cl.

13. Preparation of 3,4'-diacetoxy-5-levulinoxystilbene.

A 25 mL round bottom flask was charged with p-xylene (6 mL), Pd II catalyst (6.9 mg, 0.03 mmol), 1,3-bis-(2,6-diisopropylphenyl) imidazolinium chloride (12.9 mg, 0.03 mmol), 3-acetoxy-5-levulinoxybenzoyl chloride (0.936 g, 3 mmol), 4-acetoxystyrene (0.583 g, 3.6 mmol), and N-methyl morpholine (0.42 g, 3.6 mmol). The mixture was stirred at 120 °C for 3.5 h under nitrogen atmosphere. Then it was cooled to room temperature and EtOAc was added and filtered. The filtrate was washed with brine and dried over Na₂SO₄. Then it was filtered and purified via flash chromatography and gave the product (0.86 g, 70%) as white solid. Data are: ¹H NMR (CDCl₃, 300 MHz) δ 7.48 (m, 2H), 7.12 (m, 6H), 6.81 (t, 1H, J=1.2Hz), 2.85 (m, 4H), 2.31 (s, 1H), 2.30 (s, 1H), 2.23 (s, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ


206.4, 171.2, 169.5, 169.1, 151.5, 150.6, 140.0, 134.7, 129.8, 127.8, 127.4, 122.1, 117.14, 117.09, 114.6, 38.1, 30.0, 28.4, 21.32, 21.29.

14. Preparation of 3,4'-diacetoxy-5-hydroxystilbene.

To a solution of 3,4'-diacetoxy-5-levulinoxystilbene (0.72 g, 1.75mmol) in
5 THF (5 mL) was added a solution of Na₂SO₃ and Na₂S₂O₅ (0.26 g, 2.1 mmol and 0.1
g, 0.53 mmol) in water (5 mL). The reaction mixture was stirred for 9 h at room
temperature under N₂. Then the mixture was poured into water and extracted with
ethyl acetate three times and dried over Na₂SO₄. The solvent was removed under
reduced pressure and purified by radial chromatography to give 3,4'-diacetoxy-5-
10 hydroxystilbene (0.40 g, 73%). Data are: ¹H NMR (Aceton-d₆, 300 MHz) δ 8.66 (bs,
1H), 7.63 (m, 2H), 7.25 (m, 4H), 6.94 (t, 1H, J=1.8Hz), 6.86 (t, 1H, J=1.8Hz), 6.54
(t, 1H, J=2.1Hz), 2.26 (s, 3H), 2.25 (s, 3H); ¹³C NMR (Aceton-d₆, 75 MHz) δ 169.7,
169.6, 159.2, 153.2, 151.6, 140.4, 135.6, 129.4, 128.8, 128.3, 122.9, 111.85, 111.79,
109.4, 21.07, 21.02; HRMS (EI⁺) found 312.0996 M⁺, calcd 312.0998 for C₁₈H₁₆O₅.

The following are prepared according to Scheme 14.

Scheme 14.



5 15. Preparation of 4-levulinoxystyrene.

A flame dried flask was charged with dioxane (120 mL), 4-hydroxystyrene (3.66 g, 30 mmol), levulinic acid (6.97 g, 60 mmol), DCCl (12.39 g, 60 mmol), DMAP (300 mg). The mixture was stirred under N₂ for 3.5 h. Then the mixture was washed with water, dried over Na₂SO₄, concentrated under reduced pressure and purified by flash column to give 4-levulinoxystyrene (5.9 g, 90%). Data are: ¹H NMR (CDCl₃, 300 MHz) δ 7.39 (m, 2H), 7.04 (m, 2H), 6.68 (dd, 1H), 5.69 (d, 1H, J=17.7Hz), 5.69 (d, 1H, J=10.8Hz), 2.82 (m, 4H), 2.20 (s, 3H); ¹³C NMR (CDCl₃, 75 MHz) δ 206.5, 171.5, 150.4, 136.0, 135.5, 127.3, 121.7, 114.1, 38.0, 30.0, 28.3; HRMS (EI⁺) found 218.0955 M⁺, calcd 218.0943 for C₁₃H₁₄O₃.

16. Preparation of 3-acetoxy-4',5-dilevulinoxystilbene.

A 25 mL round bottom flask was charged with p-xylene (6 mL), Pd II catalyst (6.9 mg, 0.03 mmol), 1,3-bis-(2,6-diisopropylphenyl) imidazolinium chloride (12.9 mg, 0.03 mmol), 3-acetoxy-5-levulinoxystyrene (0.936 g, 3 mmol), 4-levulinoxystyrene (0.786 g, 3.6 mmol), and N-methyl morpholine (0.42 g, 3.6 mmol). The mixture was stirred at 120 °C for 3.5 h under nitrogen atmosphere. Then it was cooled to room temperature and EtOAc was added and filtered. The filtrate was washed with brine and dried over Na₂SO₄. Then it was filtered and purified via flash chromatography and gave the product (1.0 g, 72%) as white solid.

10 Data are: ¹H NMR (CDCl₃, 300 MHz) δ 7.47 (m, 2H), 7.11-6.92 (m, 6H), 6.81 (t, 2.1Hz), 2.89-2.80 (m, 8H), 2.29 (s, 3H), 2.224 (s, 3H), 2.221 (s, 3H); ¹³C NMR (CDCl₃, 75 MHz) δ 206.6, 206.5, 171.5, 171.2, 169.1, 151.4, 151.5, 150.6, 139.7, 134.6, 129.8, 127.8, 127.3, 122.0, 117.10, 117.05, 114.5, 38.07, 38.02, 30.01, 30.00, 28.32, 28.30, 21.3; HRMS (EI⁺) found 466.1636 M⁺, calcd 466.1628 for C₂₆H₂₆O₈.

17. Preparation of 3-acetoxy-4',5-dihydroxystilbene.


A solution of 3-acetoxy-4',5-dilevulinoxystilbene (0.47 g, 1.0 mmol) in THF (3 mL) was added a solution of Na₂SO₃ and Na₂S₂O₅ (0.30 g, 2.4 mmol and 0.11 g, 0.6 mmol) in water (3 mL). The reaction mixture was stirred for 7 h at room temperature under N₂. Then the mixture was poured into water and extracted with ethyl acetate three times and dried over Na₂SO₄. The solvent was removed under reduced pressure and purified by chromatography to give 3-acetoxy-4',5-dihydroxystilbene (0.19 g, 70%). Data are: ¹H NMR (Aceton-d₆, 300 MHz) δ 8.56 (bs, 1H), 7.45 (m, 2H), 7.04 (dd, 2H, J=48.9Hz), 6.90-6.80 (m, 4H), 6.50 (t, 1H, 2.1Hz), 2.24 (s, 1H); ¹³C NMR (Aceton-d₆, 75 MHz) δ 169.6, 159.2, 158.5, 153.3, 141.1, 129.8, 130.3, 129.0, 116.5, 111.5, 111.4, 108.8, 21.1; HRMS (EI⁺) found 270.0896 M⁺, calcd 270.0892 for C₁₆H₁₆O₄.

20

25

The following are prepared according to Scheme 15.

Scheme 15.




18. Preparation of 3,5,4'-triacetoxystilbene.

A 100 mL round bottom flask was charged with p-xylene (56 mL), $\text{Pd}(\text{OAc})_2$ (62.86 mg, 0.28 mmol), 1,3-bis-(2,6-diisopropylphenyl) imidazolium chloride (119.53 mg, 0.28 mmol), 3,5-diacetoxycarbonyl chloride (7.19 g, 28 mmol), 4-acetoxystyrene (5.35 mL, 33.6 mmol), and N-methyl morpholine (4.2 mL, 33.6 mmol). The mixture was stirred at 120°C for 3.5 h at nitrogen atmosphere. Then it was cooled to room temperature and EtOAc was added and filtered. The filtrate was washed with brine and dried over Na_2SO_4 . Then it was filtered and purified via flash chromatography (15% EtOAc/hexane, 25% EtOAc/Hexene, then 35% EtOAc/hexane) and gave resveratrol triacetate (6.95g, 70.1%) as white solid. Data are: ^1H NMR (CDCl_3 , 300 MHz) δ 7.50 (d, 2H), 7.12-6.94 (m, 6H), 6.82 (t, 1H), 2.31(s, 9H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 169.6, 169.2, 151.5, 150.6, 139.7, 134.7, 129.9, 127.9, 127.4, 122.1, 117.1, 114.6, 21.4; mp = $116-118^\circ\text{C}$; HRMS (EI^+) found 354.1118 M^+ , calcd 354.1103 for $\text{C}_{20}\text{H}_{18}\text{O}_6$; Anal. Calcd for $\text{C}_{20}\text{H}_{18}\text{O}_6$: C, 67.79, H, 5.12. Found: C, 67.93, H, 5.26.

The following are prepared according to Scheme 16.

Scheme 16.



19. Preparation of 3,5,4'-trifluorostilbene.

A 25 mL round bottom flask was charged with p-xylene (10 mL), Pd(OAc)₂ (11.3 mg, 0.05 mmol), 1,3-bis-(2,6-diisopropylphenyl) imidazolinium chloride (21.4 mg, 0.05 mmol), 3,5-difluorobenzoyl chloride (0.89g, 5 mmol), 4-fluorostyrene (0.74g, 6 mmol), and N-methyl morpholine (0.69 g, 6 mmol). The mixture was stirred at 120 °C for 3.5 h under nitrogen atmosphere. Then it was cooled to room temperature and EtOAc was added and filtered. The filtrate was washed with brine and dried over Na₂SO₄. Then it was filtered and purified via flash chromatography and gave the product (0.94g, 80%) as white solid. Data are: ¹H NMR (CDCl₃, 300 MHz) δ 7.47 (m, 2H), 7.08-6.88 (m, 6H), 6.70 (tt, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 163.52 (dd), 162.98 (d), 140.82 (t), 132.77, 130.26, 128.58 (d), 126.53, 116.03 (d), 109.20 (q), 102.98(t); ¹⁹F NMR (CDCl₃, 282 MHz) δ -31207.2 (q, 2F), -31965.6 (q, 1F); HRMS (EI⁺) found 234.0645 M⁺, calcd 234.0656 for C₁₄H₉F₃.

20. Preparation of 4'-acetoxy-3,5-difluorostilbene.

A 25 mL round bottom flask was charged with p-xylene (10 mL), Pd(OAc)₂ (11.3 mg, 0.05 mmol), 1,3-bis-(2,6-diisopropylphenyl) imidazolinium chloride (21.4 mg, 0.05 mmol), 3,5-difluorobenzoyl chloride (0.89g, 5 mmol), 4-acetoxystyrene (0.97 g, 6 mmol), and N-methyl morpholine (0.69 g, 6 mmol). The mixture was stirred at 120 °C for 3.5 h under nitrogen atmosphere. Then it was cooled to room temperature and EtOAc was added and filtered. The filtrate was washed with brine and dried over Na₂SO₄. Then it was filtered and purified via flash chromatography and gave the product (1.02g, 74%) as white solid. Data are: ¹H NMR (CDCl₃, 300 MHz) δ 7.51 (d, J= 8.4 Hz, 2H), 7.12-6.97 (m, 6H), 6.70 (t, 1H), 2.31(s, 3H).

21. Preparation of 3,5-difluoro-4'-hydroxystilbene.

To a solution of 3,5-difluoro-4'-acetoxystilbene (0.69g, 2.5mmol) in MeOH (10ml) was added 5mL 4M HCl in 1,4-dioxane. The mixture was stirred for 20 minutes at ambient temperature. After purification by flash column, 3,5-difluoro-4'-hydroxystilbene (0.51g, 88%) was obtained. Data are: ¹H NMR (CDCl₃, 300 MHz) δ 7.40 (m, 2H), 7.05-6.83 (m, 6H), 6.67 (tt, 1H), 4.88 (s, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 164.49(d), 162.47, 156.01, 141.29, 130.90, 128.53, 124.74, 115.96, 109.03 (t), 102.53 (t); ¹⁹F NMR (CDCl₃, 282 MHz) δ -31278.95 (t, J=1720.2Hz); HRMS (EI⁺) found 232.0715 M⁺, calcd 232.0700 for C₁₄H₁₀F₂O.

22. Preparation of 3,5-diacetoxy-4'-fluorostilbene.

A 25 mL round bottom flask was charged with p-xylene (10 mL), Pd(OAc)₂ (11.3 mg, 0.05 mmol), 1,3-bis-(2,6-diisopropylphenyl) imidazolinium chloride (21.4 mg, 0.05 mmol), 3,5-diacetoxybenzoyl chloride (1.28g, 5 mmol), 4-fluorostyrene (0.73 g, 6 mmol), and N-methyl morpholine (0.69 g, 6 mmol). The mixture was stirred at 120 °C for 3.5 h under nitrogen atmosphere. Then it was cooled to room temperature and EtOAc was added and filtered. The filtrate was washed with brine and dried over Na₂SO₄. Then it was filtered and purified via flash chromatography and gave the product (1.15g, 73%) as white solid. Data are: ¹H NMR (CDCl₃, 300 MHz) δ 7.45 (dd, 2H), 7.11-6.91 (m, 6H), 6.82 (t, J=1.95Hz, 1H).

23. Preparation of 4'-fluoro-3,5-dihydroxystilbene.

To a solution of 3,5-diacetoxy-4'-fluorostilbene (0.79g, 2.5mmol) in MeOH (10ml), 10mL 4M HCl in 1,4-dioxane was added. The mixture was stirred for 30 minutes at ambient temperature. After purification by flash column, 3,5-difluoro-4'-hydroxy-stilbene (0.49g, 85%) was obtained. Data are: ¹H NMR (Aceton-d₆, 300 MHz) δ 8.25 (s, 1H), 7.61 (m, 2H), 7.16-7.00 (m, 4H), 6.58 (d, J=2.4Hz), 6.31 (t, J=2.3Hz); ¹³C NMR (Aceton-d₆, 75 MHz) δ; ¹⁹F NMR (Aceton-d₆, 282 MHz) δ - 32881.41 (m); HRMS (EI⁺) found 230.0737 M⁺, calcd 230.0743 for C₁₄H₁₁FO₂.

24. Preparation of 5-benzyloxy-3-fluorophenyl bromide.

To a stirred solution of benzyl alcohol (3.86g, 20mmol) in DMA (30ml), NaH(60% dispersion in oil, 0.80g, 20mmol) was drop wisely added. After one hour, 1-bromo-3,5-difluorobenzen (3.86g, 20mmol) was added at such a rate to maintain the temperature no more than 40°C. The mixture was stirred at ambient temperature overnight. The reaction was quenched by water and extracted with EtOAc. It was purified via flash chromatography and gave the product (4.17g, 74%) as colorless oil. Data are: ¹H NMR (CDCl₃, 300 MHz) δ 7.41-7.43 (m, 5H), 6.94 (m, 1H), 6.86 (dt, 1H), 6.63 (dt, 1H), 5.02 (s, 2H); ¹³C NMR (CDCl₃, 75 MHz) δ 165.20, 161.89, 160.72, 160.56, 136.01, 128.94, 128.57, 127.77, 127.73, 122.99, 122.83, 114.58, 114.53, 112.18, 111.84, 102.29, 101.96, 70.76; ¹⁹F NMR (CDCl₃, 282 MHz) δ - 31129.40 (t, J=2583.12Hz).

25. Preparation of 5-benzyloxy-3, 4'-difluorostilbene.

A mixture of 5-benzyloxy-3-fluorophenyl Bromide (2.81g, 10mmol), Pd(OAc)₂ (0.067g, 0.3mmol), tri-*o*-tolylphosphine (0.18g, 0.6mmol), 4-fluorostyrene (1.53g, 12.5mmol), and Et₃N (1.26g, 12.5mmol) in 20ml dry DMF was stirred at 100 °C for 24 hours under N₂. The dark mixture was distributed between EtOAc and 1N HCl. The organic layer was separated and washed with water and brine, dried over Na₂SO₄, filtered, and evaporated. After flash column, the desired product (2.93g, 91%) was obtained. Data are: ¹H NMR (CDCl₃, 300 MHz) δ 7.49-7.34 (m, 7H), 7.08-6.81 (m, 6H), 6.60 (dt, 1H), 5.08 (s, 2H); ¹⁹F NMR (CDCl₃, 282 MHz) δ -31666.54 (t), -32136.54 (t). HRMS (EI⁺) found 322.1169 M⁺, calcd 322.1169 for C₂₁H₁₆F₂O.

26. Preparation of 3, 4'-difluoro-5-hydroxystilbene.

A 1 M solution of BBr₃ (9 ml, 9mmol) in CH₂Cl₂ was added to a well-stirred solution of 5-benzyloxy-3, 4'-difluorostilbene (0.97g, 3mmol) in dry CH₂Cl₂ (120 ml) at -78 °C. The mixture was warmed to -20 °C and stirred for 10 minutes. Then MeOH was added at -78 °C. The mixture was washed with saturated NaHCO₃, and water. Flash chromatography gave the 3, 4'-difluoro-5-hydroxystilbene (0.60g, 86%) as white solid. Data are: ¹H NMR (CDCl₃, 300 MHz) δ 7.46 (m, 2H), 7.08-6.73 (m, 6H), 6.48 (dt, 1H), 4.85 (s, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 165.05 (d), 161.78 (d), 157.06 (d), 140.43 (d), 133.06 (d), 129.40 (s), 128.45 (d), 127.25 (t), 115.95 (d), 109.42 (d), 105.91 (d), 102.60 (d); ¹⁹F NMR (CDCl₃, 282 MHz) δ -31703.17 (t), -32099.92 (t); HRMS (EI⁺) found 232.0714 M⁺, calcd 232.0700 for C₁₄H₁₀F₂O.

27. Preparation of 4'-acetoxy-5-benzyloxy-3-fluorostilbene.

A mixture of 5-benzyloxy-3-fluorophenyl bromide (2.81g, 10mmol), Pd(OAc)₂ (0.067g, 0.3mmol), tri-*o*-tolylphosphine (0.18g, 0.6mmol), 4-acetoxystyrene (2.03g, 12.5mmol), and Et₃N (1.26g, 12.5mmol) in 20mL dry DMF was stirred at 100 °C for 24 hours under N₂. The dark mixture was distributed between EtOAc and 1N HCl. The organic layer was separated and washed with water and brine, dried over Na₂SO₄, filtered, and evaporated. After flash column, the desired product (3.19g, 88%) was obtained. Data are: ¹H NMR (CDCl₃, 300 MHz) δ 7.51-7.34 (m, 7H), 7.11-6.82 (m, 6H), 6.60 (dt, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ

169.66, 165.64, 162.40, 160.47, 150.58, 140.07 (d), 136.59, 134.76, 129.41, 128.91, 128.41, 128.04 (d), 127.79 (d), 122.11, 109.36 (d), 105.86 (d), 101.91 (d), 70.57, 21.37; ^{19}F NMR (CDCl_3 , 282 MHz) δ -31675.70 (t); HRMS (EI^+) found 362.1302 M^+ , calcd 362.1318 for $\text{C}_{23}\text{H}_{19}\text{FO}_3$.


5 28. Preparation of 3-fluoro-5,4'-dihydroxystilbene.

A 1 M solution of BBr_3 (9 ml, 9mmol) in CH_2Cl_2 was added to a well-stirred solution of 5-benzyloxy-3-fluoro-4'-acetoxystilbene (1.09g, 3mmol) in dry CH_2Cl_2 (120 mL) at -78°C . The mixture was warmed to -20°C and stirred for 10 minutes. Then MeOH was added at -78°C . The mixture was washed with saturated NaHCO_3 ,
10 and water, dried over Na_2SO_4 , filtered and evaporated. The 4'-acetoxy-3-fluoro-5-hydroxystilbene (0.50g, 61%) was obtained as white solid. To a solution of the crude 4'-acetoxy-3-fluoro-5-hydroxystilbene (0.50g, 1.8mmol) in MeOH (7.5 ml) was added 3.7ml 4M HCl in 1,4-dioxane. The mixture was stirred for 20 minutes at ambient temperature. After purification by flash column, 3-fluoro-5,4'-
15 dihydroxystilbene (0.33g, 78%) was obtained. Data are: ^1H NMR (Aceton- d_6 , 300 MHz) δ 8.624 (s, 2H), 7.46 (m, 2H), 7.06 (q, 2H), 6.84 (m, 4H), 6.66 (dt, 1H); ^{13}C NMR (Aceton- d_6 , 75 MHz) δ 164.90 (d), 159.96 (d), 158.57, 142.02 (d), 130.84, 129.63, 129.10, 125.55 (d), 116.53, 110.28 (d), 104.44 (d), 102.00 (d); ^{19}F NMR (Aceton- d_6 , 282 MHz) δ -32429.72 (t); HRMS (EI^+) found 230.0738 M^+ , calcd
20 230.0743 for $\text{C}_{14}\text{H}_{11}\text{FO}_2$.


CLAIMS

We claim:


1. A process for preparing a compound of Formula 1A3:



- 5 where A_1 is selected from P_1 and $(CO)R_1$, A_2 is selected from P_2 and $(CO)R_2$, A_3 is selected from P_3 and $(CO)R_3$, where P_1 when present is a first alcohol protecting group, P_2 when present is a second alcohol protecting group, P_3 when present is a third alcohol protecting group, R_1 , R_2 and R_3 when present are each independently selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers
- 10 of the foregoing, comprising:
coupling a benzoyl halide compound of Formula 1A4




where X is halogen, with a 4-substituted styrene compound of Formula 1A5



- 15 in solution or in suspension of a first solvent, transition metal catalyst, N-heterocyclic carbene-type ligand and a first base.


2. A process for preparing a compound of Formula 1A3:




- 20 where A_1 is selected from P_1 and $(CO)R_1$, A_2 is selected from P_2 and $(CO)R_2$, A_3 is selected from P_3 and $(CO)R_3$, where P_1 when present is a first alcohol protecting group, P_2 when present is a second alcohol protecting group, P_3 when present is a third alcohol protecting group, R_1 , R_2 and R_3 when present are each independently

selected from alkyl with at least two carbon atoms, aryl, aralkyl, and diastereoisomers of the foregoing, comprising:

coupling a benzoyl halide compound of Formula 1A4




5 where X is halogen, with a 4-substituted styrene compound of Formula 1A5




in solution or in suspension of a first solvent, transition metal catalyst, N-heterocyclic carbene-type ligand and a first base.


10 3. A process for preparing a compound of Formula 1B



15 where Y is independently selected from NH and O, n is equal to 0, 1, 2, 3, 4 or 5, Z is independently selected from NH and O, m is equal to 0, 1, 2, 3, 4 or 5, each A and each B is independently selected from R, P₁, P₂ and P₃ and each R is independently selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing and P₁, P₂ and P₃ are protecting groups, comprising the step of coupling a benzoyl halide compound of Formula 1B2




where X is halogen, with styrene compound of Formula 1B3




20

in solution or in suspension of a first solvent, transition metal catalyst, N-heterocyclic carbene-type ligand and a first base.


4. A process for preparing a compound of Formula 1B



where Y is independently selected from NH and O, n is equal to 0, 1, 2, 3, 4 or 5, Z is independently selected from NH and O, m is equal to 0, 1, 2, 3, 4 or 5, each A and each B is independently selected from R, P₁, P₂ and P₃ and each R is independently selected from alkyl with at least two carbon atoms, aryl, aralkyl, and diastereoisomers of the foregoing and P₁, P₂ and P₃ are protecting groups, comprising the step of coupling a benzoyl halide compound of Formula 1B2




10 where X is halogen, with styrene compound of Formula 1B3




in solution or in suspension of a first solvent, transition metal catalyst, N-heterocyclic carbene-type ligand and a first base.

15 5. A process for preparing a compound of Formula I




where each Y is -O or halogen, each Z is -O or halogen, each n is independently the value of 0, 1, 2, 3, 4 or 5, each m is independently the value of 0, 1, 2, 3, 4 or 5, each A is independently selected from P_n, R or absent, each B is independently selected from H, R or absent, each V is independently selected from P_n, straight or branched alkyl of from 1 to 6 carbon atoms and cycloalkyl of from 3 to 8 carbon atoms, alkoxy, phenyl, benzyl or halogen, each W is independently selected from H, straight or branched alkyl of from 1 to 6 carbon atoms and cycloalkyl of from 3 to 8 carbon atoms, alkoxy, phenyl, benzyl or halogen, where R is selected from the group alkyl with at least one carbon atom, aryl and aralkyl, P_n is an alcohol protecting

group and diastereoisomers of the foregoing, comprising the step of coupling a benzoyl halide compound of Formula IC




where X is halogen, with a 4-substituted styrene compound of Formula IE



5

in solution or in suspension of a first solvent, transition metal catalyst, N-heterocyclic carbene-type ligand and a first base.

6. A process for preparing a compound of Formula I




10


where each Y is -O or halogen, each Z is -O or halogen, each n is independently the value of 0, 1, 2, 3, 4 or 5, each m is independently the value of 0, 1, 2, 3, 4 or 5, each A is independently selected from P_n, R or absent, each B is independently selected from H, R or absent, each V is independently selected from P_n, straight or branched alkyl of from 2 to 6 carbon atoms and cycloalkyl of from 3 to 8 carbon atoms, alkoxy, phenyl, benzyl or halogen, each W is independently selected from H, straight or branched alkyl of from 1 to 6 carbon atoms and cycloalkyl of from 3 to 8 carbon atoms, alkoxy, phenyl, benzyl or halogen, where each R is independently selected from alkyl with at least two carbon atoms, aryl and aralkyl, P_n is an alcohol protecting group and diastereoisomers of the foregoing, comprising the step of coupling a benzoyl halide compound of Formula IC

15

20




where X is halogen, with a 4-substituted styrene compound of Formula IE



in solution or in suspension of a first solvent, transition metal catalyst, N-heterocyclic carbene-type ligand and a first base.


- 5 7. A process for preparing resveratrol from compounds of Formula 1A3



where A₁ is selected from P₁ and (CO)R₁, A₂ is selected from P₂ and (CO)R₂, A₃ is selected from P₃ and (CO)R₃, where P₁ when present is a first alcohol protecting group, P₂ when present is a second alcohol protecting group, P₃ when present is a
 10 third alcohol protecting group, R₁, R₂ and R₃ when present are each independently selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing, comprising the step or steps:

- (a) reacting said compounds of Formula 1 with a first base in a first solvent; and/or,
 15 (b) reacting said compounds of Formula 1 with one or more deprotection reactants to remove said alcohol protecting groups P₁, P₂ and P₃.


8. A process for preparing resveratrol from compounds of Formula 1A3



20 where A₁ is selected from P₁ and (CO)R₁, A₂ is selected from P₂ and (CO)R₂, A₃ is selected from P₃ and (CO)R₃, P₁ when present is a first alcohol protecting group, P₂ when present is a second alcohol protecting group, P₃ when present is a third alcohol protecting group, R₁, R₂ and R₃ when present are each independently selected from alkyl with at least two carbon atoms, aryl, aralkyl, and diastereoisomers of the
 25 foregoing, comprising the step or steps:


- (a) reacting said compounds of Formula 1 with a first base in a first solvent; and/or,
- (b) reacting said compounds of Formula 1 with one or more deprotection reactants to remove said alcohol protecting groups P₁, P₂ and P₃.

5 9. A process for preparing a compound of Formula 3




pharmaceutically and cosmetically acceptable salts thereof where R₁ is selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing, comprising:

- 10 (a) esterifying a first alcohol of resorcylic acid with an acylating agent in solution or in suspension of a first solvent;
- (b) reacting a second alcohol of resorcylic acid with a first alcohol protecting agent in solution or in suspension of a second solvent,
- wherein the sequence of steps (a) and (b) are interchangeable, to give the mono-
- 15 ester, mono-protected alcohol of Formula 3B




where P₁ is a first alcohol protecting group;

- (c) halogenating said compound of Formula 3B using a halogenating agent in solution or in suspension of a third solvent to give the acid halide of
- 20 Formula 3D




where X is halogen;

- (d) coupling said compound of Formula 3E with a compound of Formula
- 3D




where P₂ is a second alcohol protecting group, with transition metal catalyst, N-heterocyclic carbene-type ligand, and a first base in solution or in suspension of a fourth solvent to yield a compound of Formula 3F;



5

(e) deprotecting said first and second alcohol protecting groups from said compound of Formula 3F to give the compound of Formula 3.

10. A process for preparing a compound of Formula 3




10

pharmaceutically and cosmetically acceptable salts thereof where R₁ is selected from alkyl with at least two carbon atoms, aryl, aralkyl, and diastereoisomers of the foregoing, comprising:

(a) esterifying a first alcohol of resorcylic acid with an acylating agent in solution or in suspension of a first solvent;

15


(b) reacting a second alcohol of resorcylic acid with a first alcohol protecting agent in solution or in suspension of a second solvent, wherein the sequence of steps (a) and (b) are interchangeable to give the mono-ester, mono-protected alcohol of Formula 3B



20


where P₁ is a first alcohol protecting group;

(c) halogenating said compound of Formula 3B using a halogenating agent in solution or in suspension of a third solvent to give the acid halide of Formula 3D




5 where X is halogen;

(d) coupling said compound of Formula 3E with a compound of Formula 3D




10 where P₂ is a second alcohol protecting group, with transition metal catalyst, N-heterocyclic carbene-type ligand, and a first base in solution or in suspension of a fourth solvent to yield a compound of Formula 3F;



(e) deprotecting said first and second alcohol protecting groups from said compound of Formula 3F to give the compound of Formula 3.


15

11. A process for preparing a compound of Formula 4




20 pharmaceutically and cosmetically acceptable salts thereof where R₁ and R₂ are independently selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing, comprising:

(a) esterifying resorcylic acid with an acylating agent in solution or in suspension of a first solvent to give the 3,5-diester of resorcylic acid of Formula 4A;




(b) halogenating said compound of Formula 4a using a halogenating agent in solution or in suspension of a second solvent to give the acid halide of Formula 4C




5

where X is halogen;

(c) coupling said compound of Formula 4C with a compound of Formula 4D




10 where P₃ is a first alcohol protecting group, with transition metal catalyst, N-heterocyclic carbene-type ligand, and a first base in solution or in suspension of a third solvent to yield a compound of Formula 4E;



(d) deprotecting said first alcohol protecting group from said compound of Formula 4E to give the compound of Formula 4.


15

12. A process for preparing a compound of Formula 4




pharmaceutically and cosmetically acceptable salts thereof where R₁ and R₂ are independently selected from alkyl with at least two carbon atoms, aryl, aralkyl, and diastereoisomers of the foregoing, comprising:

- (a) esterifying resorcylic acid with an acylating agent in solution or in suspension of a first solvent to give the 3,5-diester of resorcylic acid of Formula 4A;




- (b) halogenating said compound of Formula 4a using a halogenating agent in solution or in suspension of a second solvent to give the acid halide of Formula 4C




where X is halogen;

- (c) coupling said compound of Formula 4C with a compound of Formula




where P₃ is a first alcohol protecting group, with transition metal catalyst, N-heterocyclic carbene-type ligand, and a first base in solution or in suspension of a third solvent to yield a compound of Formula 4E;



(d) deprotecting said first alcohol protecting group from said compound of Formula 4E to give a compound of said Formula 4.

5 13. A process for preparing a compound of Formula 5




pharmaceutically and cosmetically acceptable salts thereof where R_1 and R_3 are independently selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing, comprising:

10 (a) esterifying a first alcohol of resorcylic acid with an acylating agent in solution or in suspension of a first solvent;

(b) reacting a second alcohol of resorcylic acid with a first alcohol protecting agent in solution or in suspension of a second solvent, wherein the sequence of steps (a) and (b) are interchangeable to give the mono-ester,


15 mono-protected alcohol of Formula 5B



where P_1 is a first alcohol protecting group;


(c) halogenating said compound of Formula 5B using a halogenating agent in solution or in suspension of a third solvent to give the acid halide of

20 Formula 5D




where X is halogen;

- (d) coupling said compound of Formula 5D with a compound of Formula 5E




- 5 with transition metal catalyst, N-heterocyclic carbene-type ligand, and a first base in solution or in suspension of a fourth solvent to yield a compound of Formula 5F;




- (e) deprotecting said first and second protecting groups from said compound of Formula 5F to give a compound of said Formula 5.

- 10 14. A process for preparing a compound of Formula 5




pharmaceutically and cosmetically acceptable salts thereof where R_1 and R_3 are independently selected from alkyl with at least two carbon atoms, aryl, aralkyl, and diastereoisomers of the foregoing, comprising:

- 15 (a) esterifying a first alcohol of resorcylic acid with an acylating agent in solution or in suspension of a first solvent;
- (b) reacting a second alcohol of resorcylic acid with a first alcohol protecting agent in solution or in suspension of a second solvent,
- wherein the sequence of steps (a) and (b) are interchangeable to give the mono-ester,
- 20 mono-protected alcohol of Formula 5B




where P_1 is a first alcohol protecting group;

(c) halogenating said compound of Formula 5B using a halogenating agent in solution or in suspension of a third solvent to give the acid halide of Formula 5D




5 where X is halogen;

(d) coupling said compound of Formula 5D with a compound of Formula 5E




10 with transition metal catalyst, N-heterocyclic carbene-type ligand, and a first base in solution or in suspension of a fourth solvent to yield a compound of Formula 5F;




(e) deprotecting said first and second protecting groups from said compound of Formula 5F to give the compound of Formula 5.

15 15. A process for preparing a compound of Formula 6




pharmaceutically and cosmetically acceptable salts thereof where R₃ is selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing, comprising:

20 (a) esterifying 4-hydroxy styrene with an acylating agent in solution or in suspension of a first solvent to give the 4-ester styrene of Formula 6D;




(b) reacting resorcylic acid with an alcohol protecting agent in solution or in suspension of a second solvent to give a compound of Formula 6B




where P₁ is a first alcohol protecting group and P₂ is a second alcohol protecting group;

(c) halogenating said compound of Formula 6B using a halogenating agent in solution or in suspension of a third solvent to give the acid halide of Formula 6C




where X is halogen;

(d) coupling said compound of Formula 6D with a compound of Formula 6C with transition metal catalyst, N-heterocyclic carbene-type ligand, and a first base in solution or in suspension of a fourth solvent to yield a compound of Formula 6E;




(e) deprotecting said first alcohol protecting group from said compound of Formula 6E to give the compound of Formula 6.

16. A process for preparing a compound of Formula 6




pharmaceutically and cosmetically acceptable salts thereof where R_3 is selected from alkyl with at least two carbon atoms, aryl, aralkyl, and diastereoisomers of the foregoing, comprising:

- (a) esterifying 4-hydroxy styrene with an acylating agent in solution or in suspension of a first solvent to give the 4-ester styrene of Formula 6D;




- (b) reacting resorcylic acid with an alcohol protecting agent in solution or in suspension of a second solvent to give a compound of Formula 6B




- where P_1 is a first alcohol protecting group and P_2 is a second alcohol protecting group;

- (c) halogenating said compound of Formula 6B using a halogenating agent in solution or in suspension of a third solvent to give the acid halide of Formula 6C




where X is halogen;

- (d) coupling said compound of Formula 6D with a compound of Formula 6C with transition metal catalyst, N-heterocyclic carbene-type ligand, and a first base in solution or in suspension of a fourth solvent to yield a compound of Formula 6E;




- (e) deprotecting said first alcohol protecting group from said compound of Formula 6E to give the compound of Formula 6.

17. A process for preparing a compound of Formula 7




pharmaceutically and cosmetically acceptable salts thereof where R₁, R₂ and R₃ are independently selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing, comprising:

- (a) esterifying resorcylic acid with an acylating agent in solution or in suspension of a first solvent to give the 3,5-diester of resorcylic acid of Formula 7A;




- (b) halogenating said compound of Formula 7A using a halogenating agent in solution or in suspension of a second solvent to give the acid halide of Formula 7C




where X is halogen;

- (c) coupling said compound of Formula 7C with a compound of Formula 7D



with transition metal catalyst, N-heterocyclic carbene-type ligand, and a first base in solution or in suspension of a third solvent.


18. A process for preparing a compound of Formula 7




pharmaceutically and cosmetically acceptable salts thereof where R₁, R₂ and R₃ are independently selected from alkyl with at least two carbon atoms, aryl, aralkyl, and

5 diastereoisomers of the foregoing, comprising:

- (a) esterifying resorcylic acid with an acylating agent in solution or in suspension of a first solvent to give the 3,5-diester of resorcylic acid of Formula 7A;




- 10 (b) halogenating said compound of Formula 7A using a halogenating agent in solution or in suspension of a second solvent to give the acid halide of Formula 7C



where X is halogen;


- 15 (c) coupling said compound of Formula 7C with a compound of Formula 7D



with transition metal catalyst, N-heterocyclic carbene-type ligand, and a first base in solution or in suspension of a third solvent.


19. A compound of any of Formulas 1A3, 1B, I, 3-7 prepared from any of the processes of claims 1-18.

20. A compound of Formula 9



5 where A_1 is selected from H and $(CO)R_1$, A_2 is selected from H and $(CO)R_2$, A_3 is selected from H and $(CO)R_3$, R_1 , R_2 and R_3 when present are each independently selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing.

21. A compound of Formula 9



10 where A_1 is selected from H and $(CO)R_1$, A_2 is selected from H and $(CO)R_2$, A_3 is selected from H and $(CO)R_3$, R_1 , R_2 and R_3 when present are each independently selected from alkyl with at least two carbon atoms, aryl, aralkyl, and diastereoisomers of the foregoing.

15


22. A compound selected from the group consisting essentially of:

5,4'-dihydroxy-3-propanoate stilbene, 3,5-dihydroxy-4'-propanoate stilbene, 3,4'-dihydroxy-5-propanoate stilbene, 4'-hydroxy-3,5-dipropanoate stilbene, 5-hydroxy-3,4'-dipropanoate stilbene, 3,5,4'-tripropanoate stilbene,
 20 5,4'-dihydroxy-3-butanoate stilbene, 3,5-dihydroxy-4'-butanoate stilbene, 3,4'-dihydroxy-5-butanoate stilbene, 4'-hydroxy-3,5-dibutanoate stilbene, 5-hydroxy-3,4'-dibutanoate stilbene, 3,5,4'-tributanoate stilbene, 5,4'-dihydroxy-3-pentanoate stilbene, 3,5-dihydroxy-4'-pentanoate stilbene, 3,4'-dihydroxy-5-pentanoate stilbene, 4'-hydroxy-3,5-dipentanoate stilbene, 5-hydroxy-3,4'-dipentanoate
 25 stilbene, 3,5,4'-tripentanoate stilbene, 5,4'-dihydroxy-3-hexanoate stilbene, 3,5-

dihydroxy-4'-hexanoate stilbene, 3,4'-dihydroxy-5-hexanoate stilbene, 4'-hydroxy-
 3,5-dihexanoate stilbene, 5-hydroxy-3,4'-dihexanoate stilbene, 3,5,4'-trihexanoate
 stilbene, 5,4'-dihydroxy-3-(2,4-hexadienoate) stilbene, 3,5-dihydroxy-4'-(2,4-
 hexadienoate) stilbene, 3,4'-dihydroxy-5-(2,4-hexadienoate) stilbene, 4'-hydroxy-
 3,5-(2,4-hexadienoate) stilbene, 5-hydroxy-3,4'-(2,4-hexadienoate) stilbene, 3,5,4'-
 tri-(2,4-hexanoate) stilbene, 5,4'-dihydroxy-3-dodecanoate stilbene, 3,5-dihydroxy-
 4'-dodecanoate stilbene, 3,4'-dihydroxy-5-dodecanoate stilbene, 4'-hydroxy-3,5-
 dodecanoate stilbene, 5-hydroxy-3,4'-dodecanoate stilbene, 3,5,4'-tridodecanoate
 stilbene, 5,4'-dihydroxy-3-hexadecanoate stilbene, 3,5-dihydroxy-4'-hexadecanoate
 stilbene, 3,4'-dihydroxy-5-hexadecanoate stilbene, 4'-hydroxy-3,5-hexadecanoate
 stilbene, 5-hydroxy-3,4'-hexadecanoate stilbene, 3,5,4'-trihexadecanoate stilbene,
 5,4'-dihydroxy-3-octadecanoate stilbene, 3,5-dihydroxy-4'-octadecanoate stilbene,
 3,4'-dihydroxy-5-octadecanoate stilbene, 4'-hydroxy-3,5-octadecanoate stilbene, 5-
 hydroxy-3,4'-octadecanoate stilbene, 3,5,4'-trioctadecanoate stilbene, 5,4'-
 dihydroxy-3-(9-octadecenoate) stilbene, 3,5-dihydroxy-4'-(9-octadecenoate)
 stilbene, 3,4'-dihydroxy-5-(9-octadecenoate) stilbene, 4'-hydroxy-3,5-(9-
 octadecenoate) stilbene, 5-hydroxy-3,4'-(9-octadecenoate) stilbene, 3,5,4'-tri-(9-
 octadecenoate) stilbene, 5,4'-dihydroxy-3-(9,12-octadecadienoate) stilbene, 3,5-
 dihydroxy-4'-(9,12-octadecadienoate) stilbene, 3,4'-dihydroxy-5-(9,12-
 octadecadienoate) stilbene, 4'-hydroxy-3,5-(9,12-octadecadienoate) stilbene, 5-
 hydroxy-3,4'-(9,12-octadecadienoate) stilbene, 3,5,4'-tri-(9,12-octadecadienoate)
 stilbene, 5,4'-dihydroxy-3-(6,9,12-octadecatrienoate) stilbene, 3,5-dihydroxy-4'-
 (6,9,12-octadecatrienoate) stilbene, 3,4'-dihydroxy-5-(6,9,12-octadecatrienoate)
 stilbene, 4'-hydroxy-3,5-(6,9,12-octadecatrienoate) stilbene, 5-hydroxy-3,4'-(6,9,12-
 octadecatrienoate) stilbene, 3,5,4'-tri-(6,9,12-octadecatrienoate) stilbene, 5,4'-
 dihydroxy-3-(9,12,15-octadecatrienoate) stilbene, 3,5-dihydroxy-4'-(9,12,15-
 octadecatrienoate) stilbene, 3,4'-dihydroxy-5-(9,12,15-octadecatrienoate) stilbene,
 4'-hydroxy-3,5-(9,12,15-octadecatrienoate) stilbene, 5-hydroxy-3,4'-(9,12,15-
 octadecatrienoate) stilbene, 3,5,4'-tri-(9,12,15-octadecatrienoate) stilbene, 5,4'-
 dihydroxy-3-(3,6,9-octadecatrienoate) stilbene, 3,5-dihydroxy-4'-(3,6,9-
 octadecatrienoate) stilbene, 3,4'-dihydroxy-5-(3,6,9-octadecatrienoate) stilbene, 4'-
 hydroxy-3,5,6,9-octadecatrienoate) stilbene, 5-hydroxy-3,4'-(3,6,9-
 octadecatrienoate) stilbene, 3,5,4'-tri-(3,6,9-octadecatrienoate) stilbene, 5,4'-


- dihydroxy-3-(5,8,11,14-eicosatetraenoate) stilbene, 3,5-dihydroxy-4'-(5,8,11,14-eicosatetraenoate) stilbene, 3,4'-dihydroxy-5-(5,8,11,14-eicosatetraenoate) stilbene, 4'-hydroxy-3,5-(5,8,11,14-eicosatetraenoate) stilbene, 5-hydroxy-3,4'-(5,8,11,14-eicosatetraenoate) stilbene, 3,5,4'-tri-(5,8,11,14-eicosatetraenoate) stilbene, 5,4'-
 5 dihydroxy-3-(5,8,11,14,17-eicosapentaenoate) stilbene, 3,5-dihydroxy-4'-(5,8,11,14,17-eicosapentaenoate) stilbene, 3,4'-dihydroxy-5-(5,8,11,14,17-eicosapentaenoate) stilbene, 4'-hydroxy-3,5-(5,8,11,14,17-eicosapentaenoate) stilbene, 5-hydroxy-3,4'-(5,8,11,14,17-eicosapentaenoate) stilbene, 3,5,4'-tri-(5,8,11,14,17-eicosapentaenoate) stilbene, 5,4'-dihydroxy-3-(4,7,10,13,16,19-
 10 docosaheptaenoate) stilbene, 3,5-dihydroxy-4'-(4,7,10,13,16,19-docosaheptaenoate) stilbene, 3,4'-dihydroxy-5-(4,7,10,13,16,19-docosaheptaenoate) stilbene, 4'-hydroxy-3,5-(4,7,10,13,16,19-docosaheptaenoate) stilbene, 5-hydroxy-3,4'-(4,7,10,13,16,19-docosaheptaenoate) stilbene, 3,5,4'-tri-(4,7,10,13,16,19-docosaheptaenoate) stilbene.

- 15 23. A compound of Formula 4C




where X is halogen, R₁ and R₂ are each independently selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing.

- 20 24. A compound of Formula 4C




where X is halogen, R₁ and R₂ are each independently selected from alkyl with at least two carbon atoms, aryl, aralkyl, and diastereoisomers of the foregoing.

25. A compound of Formula 1A₄



X is halogen, A₁ is selected from P₁ and (CO)R₁, A₂ is selected from P₂ and (CO)R₂, where P₁ when present is a first alcohol protecting group, P₂ when present is a
 5 second alcohol protecting group, R₁ and R₂ when present are each independently selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing.

26. A compound of Formula 1A₄




10 X is halogen, A₁ is selected from P₁ and (CO)R₁, A₂ is selected from P₂ and (CO)R₂, where P₁ when present is a first alcohol protecting group, P₂ when present is a second alcohol protecting group, R₁ and R₂ when present are each independently selected from alkyl with at least two carbon atoms, aryl, aralkyl, and diastereoisomers of the foregoing.

15 27. A compound selected from the group consisting essentially of:
 3,5-diacetoxybenzoyl chloride, 3,5-diacetoxybenzoyl bromide, 3,5-diacetoxybenzoyl
 iodide, 3,5-dipropionate benzoyl chloride, 3,5-dipropionate benzoyl bromide, 3,5-
 dipropionate benzoyl iodide, 3,5-dibutanoate benzoyl chloride, 3,5-dibutanoate
 benzoyl bromide, 3,5-dibutanoate benzoyl iodide, 3,5-dipentanoate benzoyl
 20 chloride, 3,5-dipentanoate benzoyl bromide, 3,5-dipentanoate benzoyl iodide, 3,5-
 dihexanoate benzoyl chloride, 3,5-dihexanoate benzoyl bromide, 3,5-dihexanoate
 benzoyl iodide, 3,5-dihexadienoate benzoyl chloride, 3,5-dihexadienoate benzoyl
 bromide, 3,5-dihexadienoate benzoyl iodide, 3,5-didodecanoate benzoyl chloride,
 3,5-didodecanoate benzoyl bromide, 3,5-didodecanoate benzoyl iodide, 3,5-
 25 dihexadecanoate benzoyl chloride, 3,5-dihexadecanoate benzoyl bromide, 3,5-
 dihexadecanoate benzoyl iodide, 3,5-dioctadecanoate benzoyl chloride, 3,5-

dioctadecanoate benzoyl bromide, 3,5-dioctadecanoate benzoyl iodide, 3,5-di-(9-octadecenoate) benzoyl chloride, 3,5-di-(9-octadecenoate) benzoyl bromide, 3,5-di-(9-octadecenoate) benzoyl iodide, 3,5-di-(9,12-octadecadienoate) benzoyl chloride, 3,5-di-(9,12-octadecadienoate) benzoyl bromide, 3,5-di-(9,12-octadecadienoate) benzoyl iodide, 3,5-di-(6,9,12-octadecatrienoate) benzoyl chloride, 3,5-di-(6,9,12-octadecatrienoate) benzoyl bromide, 3,5-di-(6,9,12-octadecatrienoate) benzoyl iodide, 3,5-di-(9,12,15-octadecatrienoate) benzoyl chloride, 3,5-di-(9,12,15-octadecatrienoate) benzoyl bromide, 3,5-di-(9,12,15-octadecatrienoate) benzoyl iodide, 3,5-di-(3,6,9-octadecatrienoate) benzoyl chloride, 3,5-di-(3,6,9-octadecatrienoate) benzoyl bromide, 3,5-di-(3,6,9-octadecatrienoate) benzoyl iodide, 3,5-di-(5,8,11,14-eicosatetraenoate) benzoyl chloride, 3,5-di-(5,8,11,14-eicosatetraenoate) benzoyl bromide, 3,5-di-(5,8,11,14-eicosatetraenoate) benzoyl iodide, 3,5-di-(5,8,11,14,17-eicosapentaenoate) benzoyl chloride, 3,5-di-(5,8,11,14,17-eicosapentaenoate) benzoyl bromide, 3,5-di-(5,8,11,14,17-eicosapentaenoate) benzoyl iodide, 3,5-di-(4,7,10,13,16,19-docosahexaenoate) benzoyl chloride, 3,5-di-(4,7,10,13,16,19-docosahexaenoate) benzoyl bromide, and 3,5-di-(4,7,10,13,16,19-docosahexaenoate) benzoyl iodide.


28. A compound selected from the group consisting essentially of: 3,5,-diacetoxy benzoyl chloride, 3,5,-diacetoxy benzoyl bromide, 3,5,-diacetoxy benzoyl iodide, 3-acetoxy-5-levulinoxy benzoyl chloride, 3-acetoxy-5-levulinoxy benzoyl bromide, 3-acetoxy-5-levulinoxy benzoyl iodide, 3-acetoxy-5-(methoxymethoxy) benzoyl chloride, 3-acetoxy-5-(methoxymethoxy) benzoyl bromide, 3-acetoxy-5-(methoxymethoxy) benzoyl iodide, 3-(levulinoxy)-5-(methoxymethoxy) benzoyl chloride, 3-(levulinoxy)-5-(methoxymethoxy) benzoyl bromide, 3-(levulinoxy)-5-(methoxymethoxy) benzoyl iodide, 3,5-bis(levulinoxy) benzoyl chloride, 3,5-bis(levulinoxy) benzoyl bromide, 3,5-bis(levulinoxy) benzoyl iodide, 3,5-bis(methoxymethoxy) benzoyl chloride, 3,5-bis(methoxymethoxy) benzoyl bromide, and 3,5-bis(methoxymethoxy) benzoyl iodide.

29. A compound of the formula



where each Y is independently selected from O(CO)R and halogen, n is equal to 0, 1, 2, 3, 4 or 5, Z is independently selected from O(CO)R and halogen, m is equal to 0 1, 2, 3, 4 or 5, the sum of n and m is 1 or more, each R is independently selected from alkyl with at least one carbon atom, aryl, aralkyl, and diastereoisomers of the foregoing.

30. A compound of the formula



where each Y is independently selected from O(CO)R and halogen, n is equal to 0, 1, 2, 3, 4 or 5, Z is independently selected from O(CO)R and halogen, m is equal to 0, 1, 2, 3, 4 or 5, the sum of n and m is 1 or more, each R is independently selected from alkyl with at least two carbon atoms, aryl, aralkyl, and diastereoisomers of the foregoing.

31. A compound according to claim 29 selected from the group consisting essentially of: 3,5,4'-trifluorostillbene, 4'-acetoxy-3,5-difluorostillbene, 3,5-difluoro-4'-hydroxystillbene, 3,5-diacetoxy-4'-fluorostillbene, 4'-fluoro-3,5-dihydroxystillbene, 3, 4'-difluoro-5-hydroxystillbene, and 3-fluoro-5,4'-dihydroxystillbene.

32. The processes according to any of claims 1-18 where X is Cl.

33. The processes according to any of claims 1-18 where said first base is a non-coordinating amine base.

34. The processes according to any of claims 1-18 where said first base is a non-coordinating amine base selected from the group consisting of: N,N-dimethylbenzylamine, N-methylmorpholine, and dimethylaminopyridine.

35. The processes according to any of claims 1-18 where said first base is
5 N,N-dimethylbenzylamine.

36. The processes according to any of claims 1-18 where said N-heterocyclic carbene-type ligand is selected from imidazolium, 1,3-disubstituted imidazolium, 1,3-disubstituted-4,5-dihydroimidazolium carbene-type ligands.

37. The processes according to any of claims 1-18 where said N-
10 heterocyclic carbene-type ligand is N,N'-bis-carbocycle imidazolium ligands.

38. The processes according to any of claims 1-18 where said N-heterocyclic carbene-type ligand is selected from N,N'-bis-(2,6-diisopropylphenyl) imidazolium chloride, N,N'-bis-(dimesityl) imidazolium chloride and N,N'-bis-adamantylimidazolium chloride.

39. The processes according to any of claims 1-18 where said
15 halogenating agent is selected from the group consisting essentially of: thionyl chloride, thionyl bromide, thionyl iodide, oxalyl chloride, oxalyl bromide, and oxalyl iodide.

40. The processes according to any of claims 1-18 where said
20 halogenating agent is thionyl chloride.

41. The processes according to any of claims 1-18 where said halogenating agent is oxalyl chloride.

42. The processes according to any of claims 1-18 where said acylating agent is selected from the group consisting essentially of: ester anhydrides, mixed
25 ester anhydrides, and acid halides.

43. The processes according to any of claims 1-18 where said acylating agent is acetic anhydride.

44. The processes according to any of claims 1-18 where said first protecting agent is selected from the group consisting of: MOM, and lev.

5 45. The processes according to any of claims 1-8, 11-14, and 17-19 where said second protecting agent is selected from the group consisting of: MOM, and lev.

46. The process according to any of claims 1 and 2 where said first base is selected from the group consisting essentially of: lithium hydroxide, sodium
10 hydroxide, potassium hydroxide, and pyridine.

47. The processes according to any of claims 1-18 comprising the step of irradiating said compound with ultraviolet light in an inert solvent.

48. The processes according to any of claims 1-18 where said transition metal catalyst is a Pd II catalyst selected from the group consisting of: Pd(OAc)₂,
15 PdBr₂, and PdCl₂.

49. The processes according to any of claims 1-18 where said Pd II catalyst is Pd(OAc)₂.

50. The processes according to any of claim 1-18 where the deprotection reagents are selected from the group consisting of: sodium sulfite and sodium
20 thiosulfite, aqueous pyridine at pH 6.7, NaI and TMSCl, alkaline earth metal hydroxide and a polar aprotic solvent.

51. The composition of claims 21 and 29 further comprising a therapeutically effective amount of said compound of Formula 1 in admixture with a pharmaceutically acceptable carrier.

52. The composition of claims 21 and 29 further comprising a cosmetically effective amount of said compound of Formula 1 in admixture with a cosmetically acceptable carrier are described.

53. The composition of claims 21 and 29 with mammalian enzymes
5 including SIRT1 promoting cell survival.

54. The composition of claims 21 and 29 further comprising a nutritionally effective amount of said compound of Formula 1 in admixture with a nutritionally acceptable carrier.